



Predictors of Disordered Eating in Adolescence and Young Adulthood: A Population-Based, Longitudinal Study of Females and Males in Norway

Journal:	<i>International Journal of Behavioral Development</i>
Manuscript ID:	JBD-2012-11-1840.R2
Manuscript Type:	Original Manuscript
Keywords:	disordered eating, predictors, eating problems, risk factors, gender differences, developmental phases
Abstract:	<p>We investigated longitudinal predictors for disordered eating from early adolescence to young adulthood (12-34 years) across gender and different developmental phases among Norwegian young people. Survey data from a population-based sample were collected at four time points (T) over a 13-year time span. A population-based sample of 5,679 females and males at T1 and T2, 2,745 at T3 and 2, and 718 at T4 were included in analyses, and linear regression and random intercept models were applied. In adolescence, initial disordered eating and parental overprotectiveness were more strongly related to disordered eating among females, whereas loneliness was a stronger predictor for adolescent males. Initial disordered eating during early adolescence predicted later disordered eating more strongly in late- than mid-adolescence. In young adulthood, no significant gender-specific risk factors were found. The findings provide support for both shared and specific risk factors for the developmental psychopathology of disordered eating.</p>

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Predictors of Disordered Eating in Adolescence and Young Adulthood: A Population-Based,
Longitudinal Study of Females and Males in Norway

For Peer Review

Abstract

We investigated longitudinal predictors for disordered eating from early adolescence to young adulthood (12-34 years) across gender and different developmental phases among Norwegian young people. Survey data from a population-based sample were collected at four time points (T) over a 13-year time span. A population-based sample of 5,679 females and males at T1 and T2, 2,745 at T3 and 2, and 718 at T4 were included in analyses, and linear regression and random intercept models were applied. In adolescence, initial disordered eating and parental overprotectiveness were more strongly related to disordered eating among females, whereas loneliness was a stronger predictor for adolescent males. Initial disordered eating during early adolescence predicted later disordered eating more strongly in late- than mid-adolescence. In young adulthood, no significant gender-specific risk factors were found. The findings provide support for both shared and specific risk factors for the developmental psychopathology of disordered eating.

Keywords: disordered eating, predictors, eating problems, risk factors, gender differences, developmental phases

1
2
3
4 Eating disorders, particularly anorexia nervosa, have the highest mortality of any mental illness,
5 with an estimated rate of 5.1 deaths per 1,000 person-years (Smink, van Hoeken, & Hoek,
6
7 2012). Eating disorders show high comorbidity with conditions such as major depression, anxiety
8
9 disorders, and multiple physiological complications (Fairburn & Harrison, 2003), and carry
10
11 unacceptable high personal, family, and social costs (Simon, Schmidt, & Pilling, 2005).
12
13
14

15
16 Disordered eating is usually defined as abnormal eating behaviors or symptoms of eating
17
18 disorders that are less severe than eating disorders as defined by DSM-IV-TR (American
19
20 Psychiatric Association, 2000). Despite the lower degree of severity, disordered eating is an
21
22 important predictor of eating disorders (Stice, 2002), and can also develop into full-blown eating
23
24 disorders (Shisslak, Crago, & Estes, 1995). Furthermore, disordered eating is associated with a
25
26 number of psychosocial impairments, including depressive symptoms, lowered self-esteem,
27
28 substance abuse, suicidal behaviors, and impaired functioning socially and at work (Haedt &
29
30 Keel, 2010; Mond et al., 2006; Shisslak et al., 1995). These findings show the public health
31
32 importance of both eating disorders and disordered eating.
33
34

35
36 Sociocultural, familial, personality and biological risk factors have all been hypothesized
37
38 to contribute to the development of disordered eating symptoms across development (Rosen,
39
40 1996). However, limited attention has impeded a full understanding of the impact of
41
42 developmental factors on disordered eating (Jacobi, Hayward, de Zwaan, Kraemer, & Agras,
43
44 2004; Stice, 2002). To the best of our knowledge, no longitudinal study has previously examined
45
46 whether risk factors vary across different developmental phases in both males and females.
47
48 Understanding risk across developmental phase and gender could have important implications for
49
50 the developmental psychopathology of disordered eating and the development of targeted
51
52 prevention programs. Therefore, using data from a large population-based longitudinal study in
53
54 Norway, we examined gender- and developmental phase-specific predictors of disordered eating
55
56 from early adolescence through young adulthood.
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Development occurs in sequences of phases with systematic changes in the individual's physical, cognitive, emotional, social, and behavioral development. Early patterns of adaptation in early developmental phases tend to promote adaptation in later phases, and early forms of maladaptation tend to lead to later maladaptation (Cicchetti & Rogosch, 2002; Cummings, Davies, & Campbell, 2000). This concept is the main principle in developmental psychopathology, a valuable framework to understand the development of eating problems (Rosen, 1996).

According to the developmental psychopathology perspective, examining changes in symptomatology and risk factors across development improves our understanding of causal processes (Rutter, 1988, 1989), and risk variability depending on the individual's age or development level (Rutter, 1988). Childhood risk factors may stem from a different context from that which is seen in adolescence or adulthood (Rosen, 1996). In pre-adolescence, for example, maternal dieting is a strong risk factor for excessive dieting, whereas in adolescence, appearance dissatisfaction is a strong risk factor for excessive dieting and other disordered eating behaviors (Hill, 1993).

Developmental transitions or changes may contribute to risks for physical and mental well-being, and the changes associated with the transition from childhood to early and mid-adolescence may confer a particular vulnerability to risk of psychopathology, including disordered eating (Graber, Brooks-Gunn, Paikoff, & Warren, 1994). For example, the onset of puberty and transition to adolescence leads to biologically pre-determined weight gain and appetite shifts that may result in body dissatisfaction, low self-esteem, and subsequent dieting (Jacobi et al., 2004). The transition to adolescence is also related to more autonomy and independence, which changes adolescent-parent relationships; there is increased perceived parental conflict and decreased parental support (De Goede, Branje, & Meeus, 2009). New academic and social demands placed by the transition to new schools (junior/middle schools)

1
2
3
4 may add responsibilities that stress coping capacities and compromise adjustments, which further
5
6 exacerbate to engage in disordered eating (Smolak & Levine, 1996; Smolak, Levine, & Gralen,
7
8 1993). All these changes may contribute to the development of disordered eating during
9
10 early/mid-adolescence. Late adolescence marks the transition to young adulthood, which
11
12 includes leaving home and/or attending college, reduced social support, increased dating and
13
14 sexual pressures, academic and career pressures, and identity exploration and instability (Smolak
15
16 & Levine, 1996). These stressors may play a unique role to the development of disordered eating
17
18 during late adolescence or young adulthood. The meaning of risk factors and predictors shifts
19
20 across development.
21
22
23

24 Although some risk factors for disordered eating, such as dieting, body dissatisfaction,
25
26 negative affect (mostly depressive symptoms), perfectionism, impulsivity, and substance abuse,
27
28 have been repeatedly documented in longitudinal studies (Jacobi et al., 2004; Stice, 2002), little
29
30 is known about whether these risk factors or other correlates are limited to specific
31
32 developmental phases or are active across the life span (Jacobi et al., 2004). In particular, there
33
34 are no longitudinal studies that have examined whether there are differences in the prospective
35
36 predictors of disordered eating when comparing mid-adolescence, late adolescence, and young
37
38 adulthood.
39
40
41

42 With respect to gender differences, both eating disorders and disordered eating have been
43
44 consistently found to be more common in females than males (Jacobi et al., 2004), and these
45
46 gender differences tend to escalate in adolescence, mostly after puberty. This has been attributed
47
48 to the increase in body fat that accompanies maturation, which inevitably shifts girls further away
49
50 from the thin ideal endorsed in society, with unhealthy weight regulating practices being a result
51
52 (Keel, Fulkerson, & Leon, 1997). The transition to adolescence is also associated with a greater
53
54 drop in self-esteem and an increase in depression for girls than for boys, which may add further
55
56 risks (Hankin et al., 1998; Smolak & Levine, 1996; Smolak et al., 1993). Also, the higher level of
57
58
59
60

1
2
3
4 the hormone estradiol during puberty influences body fat composition and appetite and may
5
6 trigger disordered eating behaviors among girls (Klump, Keel, Sisk, & Burt, 2010).
7
8

9 Differences in gender roles and expectations, particularly for thinness and attractive body
10 appearance, are also important factors to explain why eating disorders are more prevalent in
11 females than in males (Culbert, Racine, & Klump, 2011; Murnen & Smolak, 1997; Smolak &
12 Striegel-Moore, 1996). According to gender socialization theory, the social environment is an
13 important channel for the modeling of gender roles, where women are represented as being
14 dependent and lacking autonomy and competency (Worell & Todd, 1996). These negative
15 aspects of gendered social roles may lead to lowered self-worth, increasing self-concept
16 instability, and increasing reliance on others for approval of physical attractiveness, which may
17 in turn stress adolescent girls and women to attain the ideal of thinness and engage in disordered
18 eating behaviors.
19
20
21
22
23
24
25
26
27
28
29

30
31 Similarly, self-objectification theory offers an important explanation for gender
32 differences in disordered eating. This theory posits that girls and women are socialized to view
33 and treat their bodies as objects and to be preoccupied with the thin ideal, which in turn leads to
34 body shame or raises concerns about their physical appearance, and consequently, to disordered
35 eating, regardless of their level of body dissatisfaction (Fredrickson & Roberts, 1997;
36 Fredrickson, Roberts, Noll, Quinn, & Twenge, 1998).
37
38
39
40
41
42
43

44 However, gender differences on the level of disordered eating may not necessarily imply
45 differences between females and males in how strongly risk factors are associated with
46 disordered eating. There are similarities in the factors associated with disordered eating in
47 adolescent boys and girls (Ricciardelli & McCabe, 2004). These factors include body mass index
48 (BMI), negative affect, self-esteem, perfectionism, drug use and perceived pressure to lose
49 weight from parents and peers. With the exception of BMI, most of these correlates have only
50 been found in cross-sectional studies (Ricciardelli & McCabe, 2004).
51
52
53
54
55
56
57
58
59
60

1
2
3
4 On the other hand, recent longitudinal studies suggest that low self-esteem, perfectionism,
5
6 body dissatisfaction, and high BMI may be more prominent risk factors for adolescent girls than
7
8 boys (Beato-Fernandez, Rodriguez-Cano, Belmonte-Llario, & Martinez-Delgado, 2004; Ferreiro,
9
10 Seoane, & Senra, 2011; McCabe & Ricciardelli, 2006), whereas symptoms of depression have
11
12 been reported to be more prominent for adolescent boys (McCabe & Ricciardelli, 2006).
13
14

15
16 Nevertheless, most studies did not examine whether the gender differences were statistically
17
18 different for males and females. Moreover, research in this area has typically used small samples
19
20 with differing participant characteristics and study methodologies, which may further explain the
21
22 inconsistent findings across studies.
23

24
25 Examining whether risk factors for disordered eating differ by gender or developmental phase
26
27 may increase our understanding of the developmental psychopathology of disordered eating and
28
29 open the possibility of designing targeted preventive interventions for disordered eating on the
30
31 basis of early identifiable characteristics. For this reason, we aim to: (1) determine the
32
33 longitudinal relationship of predictors for disordered eating by examining these associations in
34
35 mid-adolescence, late adolescence, and young adulthood, for both males and females, and (2)
36
37 examine whether gender and developmental phases moderate the relationships between
38
39 predictors and disordered eating. This study includes putative bio-psychosocial risk factors of
40
41 disordered eating that are documented in the literature, such as appearance dissatisfaction, BMI,
42
43 pubertal development, negative affectivity, self-worth, and substance use behaviors (Jacobi et al.,
44
45 2004; McCabe & Ricciardelli, 2004; Stice, 2002); loneliness (Levine, 2012); self-concept
46
47 instability (Kansi, Wichstrom, & Bergman, 2003); and parental bonding styles (Tata, Fox, &
48
49 Cooper, 2001).
50
51
52
53
54
55
56
57
58
59
60

Method

Procedure and Participants

Data were analyzed from the Norwegian longitudinal study “Young in Norway”, which was conducted at four time points: 1992 (T1), 1994 (T2), 1999 (T3), and 2005 (T4). The initial sample at T1 was composed of 12,655 students in grades 7 to 12 (12 to 20 years of age) at 67 representative schools in Norway, with each grade being equally represented. Every school in the country was included in the register from which the schools were selected, and the sample was stratified according to geographical region and school size, which in Norway is closely related to the degree of urbanization. Each school’s sampling probability was proportional to the number of students at the school, thus providing an equal probability of selection for each student. The response rate at T1 was 97% (N = 12,287).

In 1994, three of the participating schools at T1 were not part of the follow-up study (T2; age 14 to 22), and at another school, a burglary in the school’s archives resulted in the loss of the project’s identification records. In total, 9,679 students at 63 schools were eligible to complete the T2 questionnaire. Since a considerable proportion of the students had completed their three-year track at the junior or senior high school that they were attending at T1, the subjects who were no longer at the same school at T2 received the questionnaire by mail. For this group, the response rate was 68% (N = 3,783), whereas those at their original schools had a response rate of 92% (N = 4,187). The overall response rate at T2 was 79%.

At T3, only students who completed the questionnaire in school at T2 (N = 3,844) were followed up due to a comparatively lower response rate among those receiving the questionnaire by mail. As such, those who responded by mail at T2 (N = 3,783) were not included in the follow up at T3 and T4. Since the survey was originally planned as a two-wave study, informed consent had to be obtained again at T2. Out of the total number of consenting individuals at T2 (N =

3,507, 91.2%), 2,923 (83.8%) responded to the questionnaire that they received by mail at T3 (age 19 to 28), representing an overall response rate of 68%.

In 2005 (T4), all persons who had consented to the follow-up at T2 were again invited to participate (age 25 to 34). In total, 2,890 of 3,507 potential participants, or 82.4%, completed the questionnaire, resulting in an overall response rate of 67%.

In this study, after excluding participants over 17 years of age at T1 ($N = 3,835$), and those with invalid data for various reasons, the actual sample sizes were 5,679 at T1 and T2, 2,745 at T3, and 2,718 at T4 (see Figure 1 for a graphical representation of the selected samples). Moreover, to examine longitudinal associations between predictors and disordered eating during young adulthood (i.e., combined T3 and T4 samples), participants who responded at both time points were included ($N = 2,164$).

Even though a large proportion of the sample that did not respond to the questionnaires at T3 and T4 were planned non-responders, analyses were conducted to explore the potential impact of variables on attrition. More specifically, we performed a multiple logistic regression to investigate whether variables at T1 predicted dropout at T2, T3, or T4. The results of these analyses revealed that older age, male gender, more occasions of alcohol intoxication over the past year, and higher perceived parental overprotectiveness significantly predicted higher odds for dropout at T2, T3, or T4 ($p < 0.05$). Lower scores for parental care and loneliness also predicted dropout ($p < 0.05$).

FIGURE 1 ABOUT HERE

Measures

All predictors and outcomes were based on adolescent self-report.

Outcome Variable. Disordered eating was assessed by the Eating Attitudes Test-12 (EAT-12), an instrument designed to measure eating problems and concerns related to dieting, bulimia, and food preoccupation and oral control (Garner, Olmsted, Bohr, & Garfinkel, 1982;

1
2
3
4 Lavik, Clausen, & Pedersen, 1991). The response alternatives range from 1 (“never”) to 4
5 (“always”). EAT-12 was constructed from the 26-item version of the EAT by selecting the items
6
7 with the best psychometric properties (Garner et al., 1982; Lavik et al., 1991). Mean scores were
8
9 calculated, with high scores reflecting high levels of disordered eating. The scale showed a good
10
11 internal consistency (Cronbach’s α) at each survey point: 0.73, 0.77, 0.77 and 0.75 at T1, T2, T3,
12
13 and T4, respectively.
14
15
16

17
18 **Predictor Variables.** Appearance satisfaction was assessed by the Body Areas
19
20 Satisfaction Scale (BASS) (Brown, Cash, & Lewis, 1989), which rates respondents’ level of
21
22 satisfaction with seven items or body areas: face, lower torso, mid-torso, upper torso, muscle
23
24 tone, weight and height, with the response options varying from 1 (“very dissatisfied”) to 5
25
26 (“very satisfied”). A mean score was computed, with high scores indicating a high level of
27
28 satisfaction, and the scale demonstrated a good internal consistency at each survey point with $\alpha =$
29
30 0.80, 0.81, 0.82, and 0.82 at T1, T2, T3, and T4, respectively.
31
32

33
34 General self-worth was measured using the Global Self-Worth Subscale from a revised
35
36 version of the Harter’s Self-Perception Profile for Adolescents (Harter, 1988; Wichstrom, 1995).
37
38 Five items assess how an adolescent views himself/herself, with the response options ranging
39
40 from 1 (“corresponds very poorly”) to 4 (“corresponds very well”). Higher mean scores reflect
41
42 high self-worth, and the scale had an acceptable internal consistency on all occasions: 0.70, 0.75,
43
44 0.75, and 0.78 at T1, T2, T3, and T4, respectively.
45
46

47
48 Negative affectivity, including symptoms of depression and anxiety was measured with a
49
50 12-item short version of the Hopkins Symptom Checklist (Derogatis, Lipman, Rickels,
51
52 Uhlenhuth, & Covi, 1974). Using a response scale ranging from 1 to 4, participants were asked to
53
54 restrict their ratings to the preceding week. Mean scores were calculated, with high scores
55
56 indicating high levels of negative affectivity, and the scale revealed a satisfactory internal
57
58
59
60

1
2
3
4 consistency on all occasions, with α values of 0.87, 0.88, 0.89, and 0.89 at T1, T2, T3, and T4,
5
6 respectively.
7

8
9 Loneliness was measured by a five-item version of the UCLA Loneliness Scale (Russell,
10
11 Peplau, & Cutrona, 1980), with each item having response options ranging from 1 (“never”) to 4
12
13 (“often”). A higher mean score reflects greater loneliness, and the five-item scale exhibited a
14
15 somewhat low internal consistency at T1 ($\alpha = 0.65$), whereas the α values were acceptable at the
16
17 remaining three time points, with $\alpha = 0.72$, 0.76, and 0.78 at T2, T3, and T4, respectively.
18
19

20
21 Alcohol intoxication was measured by asking participants to indicate how often they had
22
23 “drunk so much that you felt clearly intoxicated” during the preceding 12 months, and illicit drug
24
25 use was also assessed by asking about the frequency of illicit drug use over the past 12 months.
26
27 For both measures, the response scale ranged from 1 (“never”) to 6 (“more than 50 times”). High
28
29 mean scores indicate a high level of alcohol and illicit drug use.
30

31
32 Self-concept instability was measured using a revised version of Rosenberg’s Stability of
33
34 Self Scale (Rosenberg, 1986), which has four items, each with a response scale ranging from 1 to
35
36 4. High scores indicate an unstable perception of self (Alsaker & Olweus, 1986; Rosenberg,
37
38 1986), and the internal consistency was satisfactory at each survey point: 0.81, 0.86, 0.88, and
39
40 0.89 at T1, T2, T3, and T4, respectively.
41

42
43 Pubertal timing was measured by the Pubertal Development Scale (Petersen, Crockett,
44
45 Richards, & Boxer, 1988), which assesses the development of five pubertal indices and uses a
46
47 four-point scale ranging from “not begun” to “development completed.” Three of the indices are
48
49 gender neutral, and two are gender-specific; the indices for both boys and girls were pubic hair,
50
51 growth spurt, and skin changes. In addition, boys rated their development of facial hair and voice
52
53 change, and girls rated their breast development and whether they had reached menarche.
54
55 Pubertal timing was only assessed at T1. A higher mean score indicated early pubertal timing,
56
57
58
59
60

with the scales for both genders demonstrating a good internal consistency: 0.92 for girls and 0.90 for boys at T1.

Parental bonding styles were assessed by use of a short version of the Parental Bonding Instrument (Parker, Tupling, & Brown, 1979), which measures the emotional relationship between participants and parents by focusing on two dimensions, parental care and parental overprotection. Each dimension consists of five items and has a response scale ranging from 1 (“very like”) to 4 (“very unlike”). High mean scores on the care subscale indicate a parent–child relationship based on emotional warmth, closeness, and empathy, whereas high scores on the overprotection subscale suggest parental obstruction of independent behavior, as well as parental control and parental intrusion (Parker et al., 1979). Adolescents reported on the parental bonding styles only at T1, and the internal consistency coefficients for parental care and overprotection scales were 0.70 and 0.71 at T1, respectively.

Body mass index (BMI, kg/m^2) was computed from self-reported measures of height and weight, as self-reported BMI has been demonstrated to be a valid measure of actual BMI (Goodman, Hinden, & Khandelwal, 2000). Age and gender was recorded in all surveys; male was coded as “0” and female as “1.”

Statistical Analysis

First, we computed descriptive statistics (means, standard deviations) of all variables included in the study. To obtain figures for different developmental phases, we divided the study population into two different cohorts at T1 and T2: a younger cohort, which was defined to be in early adolescence at T1 (age 12 to 14) and mid-adolescence at T2 (14-16 years), and an older cohort being in mid-adolescence at T1 (age 15 to 17) and in late adolescence at T2 (age 17 to 19) (see Figure 1). Second, we applied linear regression models to examine the effects of predictors on disordered eating for both the younger and older cohort and both genders separately. To examine longitudinal predictors of disordered eating in mid-adolescence, disordered eating

1
2
3
4 scores at T2 for the younger cohort were regressed on predictors measured at T1. The same
5
6 analyses were conducted in the older cohort to examine predictors of disordered eating in late
7
8 adolescence. Third, since the young adulthood period was represented by two time points (T3
9
10 and T4), disordered eating scores from both time points were simultaneously included in a
11
12 regression model as an outcome variable, whereas variables at T2 were included as predictors. A
13
14 linear random intercept model was used for these regression analyses, taking the dependency of
15
16 the disordered eating scores at T3 and T4 into account (Rabe-Hesketh & Skrondal, 2008). This
17
18 model therefore provides regression estimates comparable to usual regression analyses, with an
19
20 additional estimate of how time point (T3 versus T4) influences disordered eating.
21
22
23

24
25 In all regression and linear random intercept models, initial disordered eating scores were
26
27 included as a covariate, thereby examining how predictors were related to *changes* in disordered
28
29 eating. This analytical approach helps in establishing temporal precedence between the outcome
30
31 (i.e., disordered eating) and the predictors. To test for differences in the regression coefficients
32
33 between: (i) females and males, and (ii) mid- and late adolescence, two-way interaction terms
34
35 were computed and included together with the main effects in regression analyses. Similarly, to
36
37 examine the possibility for regression coefficients to vary according to a combination of gender
38
39 and developmental phase, three-ways interactions were tested as well. As described by Dawson
40
41 and Richter (2006), the delta method was used to obtain information on the direction of the three-
42
43 ways interaction terms when such effects were found. This method compares differences among
44
45 regression coefficients that computed from the three-way interaction analyses –
46
47 gender*developmental phase*predictor (Dawson & Richter, 2006). We used a similar modeling
48
49 approach for both main effects and interaction analyses, first assessing the effect of single
50
51 predictor and then conducting multiple regression among significant predictors ($p < .05$).
52
53
54
55 Multicollinearity was examined using the Variance Inflation Factor (VIF), which in this study
56
57
58
59
60

was less than 3 for all variables, indicating few problems with multicollinearity. We used Stata SE/11 for Windows for all analyses.

Results

Descriptive Summary

Table 1 presents the means and standard deviations for all study variables in adolescence and young adulthood phases separately for females and males. Females had higher scores for disordered eating, BMI, negative affectivity, and self-concept instability, and lower scores for appearance satisfaction and self-worth than males across all developmental phases. Males had higher scores on illicit drug use and alcohol intoxication compared to females.

TABLE 1 ABOUT HERE

Predictors in Mid- and Late Adolescence

We examined how potential risk factors predicted disordered eating prospectively at mid- and late adolescence by means of linear regression analyses for both girls and boys (see Table 2). Measurements at T1 were used as independent variables and disordered eating at T2 as an outcome. In a first step, we conducted one regression analysis for each predictor while controlling for T1 disordered eating and age (see the results in Table 2 denoted 'β' models). We then performed multiple regression analyses in which all predictors with initial significant effects ($p < .05$) were simultaneously included (see results in Table 2 denoted 'β+' models). Similarly, we first tested the interaction terms of each predictor in regression models ('β') while controlling for main effects, T1 disordered eating and age. Next, the interaction effects of those significant predictors were examined one by one in multiple regression models ('β+') (interaction results not shown in Table 2). Findings from both regression models ('β' and 'β+') showed that initial disordered eating was significantly positively associated with disordered eating scores at T2 across developmental phases and genders. Furthermore, the two-way interaction terms revealed that the association between initial disordered eating and later disordered eating was stronger for

1
2
3
4 girls than for boys ($\beta = .31, t = 6.59, p < .001$ and $\beta_{+} = .29, t = 6.11, p < .001$), and stronger in late
5
6 adolescence than mid-adolescence ($\beta = .31, t = 6.64, p < .001$ and $\beta_{+} = .32, t = 5.77, p < .001$).

7
8
9 The three-way interaction term was not significant ($p > .05$).

10
11 BMI was positively related to disordered eating in all multiple regression analyses. No
12
13 significant interaction terms were detected for BMI, indicating that there were no significant
14
15 differences in the regression coefficients between the four groups. Appearance satisfaction and
16
17 self-worth were significantly related to lower disordered eating scores in all groups in the initial
18
19 analyses. However, in multiple regressions, appearance satisfaction in mid-adolescent girls and
20
21 boys, as well as self-worth in adolescent boys and late adolescent girls, failed to reach statistical
22
23 significance. Additionally, no significant interaction terms were found for these two variables.

24
25
26
27 Negative affect predicted a higher disordered eating score among boys and girls, but for
28
29 girls the significant association in late adolescence disappeared in multiple regression. The
30
31 interaction analyses in the initial “ β ” models confirmed boys’ regression coefficients for negative
32
33 affect to be significantly larger than those for girls ($\beta = -.09, t = -2.17, p < .05$). However, this
34
35 interaction effect did not retain statistical significance in multiple regressions. Loneliness
36
37 significantly predicted higher disordered eating scores in mid- and late adolescent boys, but only
38
39 the association in late adolescence remained significant in multiple regressions. The interaction
40
41 analyses also showed that loneliness had a stronger significant association with disordered eating
42
43 in boys than in girls ($\beta = -.11, t = -2.85, p < .01$ and $\beta_{+} = -.08, t = -2.41, p < .05$).

44
45
46
47 Alcohol intoxication had no significant associations in any group, while illicit drug use
48
49 significantly predicted a higher disordered eating in mid-adolescent boys and a lower disordered
50
51 eating in mid-adolescent girls, although these associations failed to reach statistical significance
52
53 in multiple regression models. Also, a significant three-way interaction term ($\beta = .32, t = 3.23, p <$
54
55 $.01$) and the post-hoc tests exhibited a stronger positive association between illicit drug use and
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

disordered eating among mid-adolescent boys (M slope = .11, $z = 2.80$, $p < .01$) compared to all other groups. Again, this significant interaction was not observed in multiple regression analyses.

Self-concept instability significantly predicted higher disordered eating scores in adolescent boys and late adolescent girls but only remained significant in late adolescent girls in multiple regression models. Moreover, a significant three-way interaction term ($\beta = .18$, $t = 2.07$, $p < .05$) and the post-hoc tests showed self-concept instability to have a stronger significant positive association among late adolescent girls (M slope = .06, $z = 4.65$, $p < .05$) compared to mid-adolescent girls and late adolescent boys. This interaction term did not retain statistical significance in multiple regression analyses.

Early pubertal development was significantly related to disordered eating among boys in late adolescence but not to the three other groups. Since the pubertal development measurement was gender-specific, we did not assess its interaction effect.

Concerning parental bonding styles, perceived parental overprotectiveness was significantly related to higher disordered eating scores among mid-adolescent boys and adolescent girls but only remained statistically significant in adolescent girls in multiple regression analysis. Its interaction term with gender became significant in multiple regression ($\beta = .09$, $t = -2.42$, $p < .05$), indicating a stronger association between parental overprotectiveness and disordered eating in adolescent girls than in boys; no other significant interaction terms were found for parental overprotectiveness. Lastly, a significant relationship between parental care and a lower disordered eating was found among mid-adolescent boys in the “ β ” model and in three-ways interaction analyses ($\beta = -.24$, $t = -2.12$, $p < .05$), although both associations did not retain statistical significance in multiple regression models.

TABLE 2 ABOUT HERE

Predictors during Young Adulthood

To estimate the effect of predictors beyond the adolescent phases, we developed random intercept models with disordered eating scores at T3 and T4 as outcome variables and measures at T2 as predictors. Table 3 shows the summarized results. Similar to the approach used in Table 2, we first conducted random intercept models for each predictor while controlling for T2 disordered eating and age (see results in Table 3 denoted ' β ') and then selected significant predictors ($p < 0.05$) for multiple random intercept models (see results in Table 3 denoted ' $\beta+$ '). The same approach also applied to examine the interaction terms in both regression models (" β " and " $\beta+$ " models). However, since both time periods (i.e., T3 and T4) represent young adulthood, we only performed two-way interaction analyses of gender (interaction results not shown in Table 3).

For both genders, disordered eating and self-concept instability at T2 were significantly positively associated with disordered eating in young adulthood. For these factors, no significant interaction terms with gender were found ($p > .05$). Appearance satisfaction significantly predicted less disordered eating in both genders, but only remained a significant predictor in young adult women in multiple regression. Although its interaction term showed a stronger association with a lower disordered eating in young adult women than young adult men ($\beta = -.04$, $z = -2.21$, $p < .05$), it was no longer statistically significant in a multiple regression analysis.

Negative affectivity significantly predicted higher disordered eating scores in both genders but only remained significant in young adult men in the multiple regression model. Its interaction effect also demonstrated a stronger association in young adult men than young adult women ($\beta = -.24$, $z = -2.12$, $p < .05$) but disappeared in the multiple regression analysis. Loneliness in both genders and parental care in young adult men showed significant associations to disordered eating in the initial models, but such relationships were not found in multiple regression models. The regression coefficients for the changes of disordered eating scores

1
2
3
4 between the survey time points (T3 and T4) were significantly positive in young adult men and
5
6 negative in young adult women, thereby revealing an increase in disordered eating score in
7
8 young adult men over time, while decreasing over time in young adult women. The interaction
9
10 term was significant ($\beta = -.12, z = -8.07, p < .001$ and $\beta+ = -.12, z = -7.74, p < .001$), indicating a
11
12 larger decrease in disordered eating score in young adult women than young adult men over time.
13
14 No other significant predictors and interaction terms were found in the regression models (“ β ”
15
16 and “ $\beta+$ ” models).
17
18

19
20 TABLE 3 ABOUT HERE
21

22 Discussion

23
24 The main aim of this population-based longitudinal study was to examine whether
25
26 predictors of disordered eating differed according to gender and developmental phases. The study
27
28 documented both shared and specific risk factors across gender and developmental phases. More
29
30 specifically, initial disordered eating and parental overprotectiveness predicted disordered eating
31
32 more strongly in adolescent girls than in adolescent boys. In contrast, loneliness predicted
33
34 disordered eating in adolescent boys more strongly than in adolescent girls. Furthermore, initial
35
36 disordered eating was a stronger predictor in late rather than in mid-adolescence. No other
37
38 associations between other risk factors and disorder eating differed significantly according to
39
40 gender or developmental phase in multiple regression models.
41
42
43

44 The stronger association between initial disordered eating and later disordered eating in
45
46 adolescent girls than boys could suggest a higher stability of disordered eating in females.
47
48 Although a few prior longitudinal studies assessing eating problems included female and male
49
50 participants, none of these studies tested for possible gender difference in the stability of eating
51
52 problems (Kotler, Cohen, Davies, Pine, & Walsh, 2001; Leon, Fulkerson, Perry, Keel, & Klump,
53
54 1999; Neumark-Sztainer, Wall, Larson, Eisenberg, & Loth, 2011). Greater stability of disordered
55
56 eating in adolescent girls may be due to social messages about the importance of thinness for
57
58
59
60

1
2
3
4 girls and greater susceptibility to peer and media pressures on girls to be thin in line with gender
5
6 socialization theory (Murnen & Smolak, 1997; Smolak & Levine, 1996). In addition, the rise in
7
8 girls' body fat during puberty (Keel et al., 1997), and the role of estradiol for body fat
9
10 composition and appetite (Klump et al., 2010) may explain the increased stability of disordered
11
12 eating in adolescent girls.
13
14

15
16 Perceived parental overprotectiveness was also stronger risk factor for adolescent girls
17
18 than boys. This stronger association may be due to the closer attachment that they have with their
19
20 parents than boys have, especially with their mothers (Gilligan, 1982; Shomaker & Furman,
21
22 2009). Adolescent girls often place more value on the quality of their family relationships
23
24 (Tomori, Zalar, & Plesnicar, 2000). In addition, other studies found that daughters gave their
25
26 parents higher scores for overprotectiveness than sons did (Murphy, Brewin, & Silka, 1997), with
27
28 overprotectiveness being significantly associated with disordered eating behaviors in young adult
29
30 females though not in young adult males (Tata et al., 2001). In general, higher parental
31
32 protectiveness could result in family conflict and stressful experiences, and disordered eating
33
34 behaviors could be a response to these negative affective states and vice versa, particularly for
35
36 the development of binge eating among females (Striegel-Moore et al., 2005). Hence, the exact
37
38 mechanism of this association should be verified in future prospective studies.
39
40
41

42
43 Loneliness appeared as a male-specific risk factor during adolescence. This finding is in
44
45 contrast to earlier studies reporting that females are more likely to report disordered eating
46
47 symptoms in response to loneliness than males are (Levine, 2012). This could be related to the
48
49 degree of sensitivity towards social relationships, as adolescent girls are found to be more
50
51 sensitive to the status of their peer relationships and friendships than adolescent boys (Rose &
52
53 Rudolph, 2006). However, there is generally a scarcity of research on the association between
54
55 loneliness and eating problems, particularly among males. Further research is recommended to
56
57 confirm our finding.
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

At the same time, even though negative affectivity was not a statistically significant predictor in multiple regression models, it consistently predicted disordered eating among adolescent boys and young adult men. These findings may suggest that disordered eating in males might be associated with an increased vulnerability to psychological co-morbidities as compared to females. Similar results have been found in one cross-sectional study that showed that binge eating and bulimic symptoms were associated with psychological problems to a larger degree in adolescent boys than in girls (McCabe & Vincent, 2003).

As for development-specific risk factors, only initial disordered eating exhibited a significantly stronger association in late than in mid-adolescence. This greater stability in late adolescence may be related to a heightened emotional and social distress. Since this phase is marked as the end of adolescence, with graduating high school and starting college, late adolescents may experience reduced social support, different standards of attractiveness, increased dating and sexual pressures, and intensified academic and peer pressures (Smolak & Levine, 1996). These changes and experiences are possibly stronger triggers for disordered eating behaviors in late adolescence than in earlier phases.

There were also risk factors that predicted disordered eating across gender and/or developmental phases in multiple regression analyses (significant main effects). Risk factors that predicted disordered eating in both genders include BMI in adolescence, appearance satisfaction in late adolescence, and self-concept instability in young adulthood. These findings suggest more similarities than differences among the risk factors of disordered eating between males and females in adolescence and young adulthood. The results are consistent with past research among adolescent boys and girls (Leon et al., 1999; Ricciardelli & McCabe, 2004). Our study further provides support for such similarity in young adulthood. Besides, significant main effects include a higher BMI in both mid and late adolescent girls and boys, appearance satisfaction in late adolescent girls and boys and in young adult women, self-worth in mid-adolescent girls, negative

1
2
3
4 affectivity in all developmental phases in males, and self-concept instability in mid- and late
5
6 adolescent girls and in young adults of both genders. With regard to a higher BMI, both
7
8 significant and non-significant results have been reported for adolescent and adult female
9
10 samples (Stice, 2002) as well as adolescent male samples (Ricciardelli & McCabe, 2004).
11
12 Regardless of age (Stice, 2002), body dissatisfaction has been mentioned as one of the most
13
14 consistent and robust risk factors for eating pathology in females but not for adolescent males
15
16 (Ricciardelli & McCabe, 2004). Prospective studies have reported mixed findings for negative
17
18 affectivity: Both significant (Leon et al., 1999) and non-significant (Stice, Akutagawa, Gaggan, &
19
20 Agras, 2000) relationships between negative affect and disordered eating were documented.
21
22 Individuals with an unstable perception of themselves, particularly in conjunction with low self-
23
24 worth (Kansi et al., 2003), could be more vulnerable to external information about how others
25
26 perceive their appearance (Campbell, 1990). This may lead to negative emotions and the
27
28 subsequent development of disordered eating. In general, some of divergent findings could be
29
30 due to the use of different assessment measures for disordered eating and predictors and
31
32 differences in methodological approaches, follow-up periods, and characteristics of sample
33
34 populations.
35
36
37
38
39

40 A main strength of the study is its longitudinal nature, which allows for a delineation of
41
42 the temporal order between predictors and outcome by controlling for initial disordered eating
43
44 symptoms. To the best of our knowledge, this is the first longitudinal study to explore the
45
46 prospective effects of predictors of disordered eating at different developmental phases in a large
47
48 population sample of adolescent and young adult males and females.
49
50

51 Nonetheless, the study also has its limitations. First, we used a general measure for
52
53 disordered eating, though predictors could vary across different types of disordered eating such
54
55 as dieting, bulimic symptoms, symptoms of anorexia nervosa, and binge eating. Although EAT-
56
57 12 has three correlated subscales, these subscales exhibit a rather low reliability, particularly in
58
59
60

1
2
3
4 relation to the Oral Control factor (Engelsen & Hagtvet, 1999). Moreover, since eating disorder
5 not otherwise specified (EDNOS) is the most common type of eating disorder (Wilfley, Bishop,
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

relation to the Oral Control factor (Engelsen & Hagtvet, 1999). Moreover, since eating disorder not otherwise specified (EDNOS) is the most common type of eating disorder (Wilfley, Bishop, Wilson, & Agras, 2007), it further justifies the use of global disordered eating as an outcome in the study. Second, even though it has been argued that full-syndrome eating disorders may not qualitatively differ from sub-threshold presentations (Stice, 2002; Stice, Killen, Hayward, & Taylor, 1998), it remains to be seen whether the findings from this study will be supported in studies examining the predictors of full-threshold eating disorders. Third, although many predictors displayed some significant effects, their regression coefficients were below 0.2, except for the effects of initial disordered eating, thus suggesting that small effect sizes should be considered when interpreting the findings. Fourth, although we established a factor's status as a risk factor due to precedence (Jacobi et al., 2004), this study cannot determine whether these risk factors were causal. Fifth, the larger proportion of lost follow-up could be a source of bias, especially since the attrition was significantly associated with participants' characteristics and behaviors, such as older age, male gender, higher perceived parental overprotectiveness, and lower scores for parental care and loneliness. These factors may under- or over-estimate the frequency and the prevalence of disordered eating and thus potentially bias regression estimates. Sixth, although age demarcations are often used for measuring developmental phases, these can be arbitrary and challenging given the variation in physical, social, and cognitive development. Finally, the larger time lag between measurements during adolescence and young adulthood could attenuate regression estimates of predictors on disordered eating.

Overall, the findings of this study may provide implications for selective prevention and intervention strategies tailored to the concerns of specific gender and developmental phases as well as universal strategies applicable across gender and adolescence and young adulthood. Since a higher BMI was found to be the most consistent predictor for disordered eating across gender and phases of adolescence, preventing actual or perceived overweight could be a natural target

1
2
3
4 for universal prevention programs. Other significant risk factors such as early disordered eating,
5
6 loneliness, and negative affectivity, which were revealed as specific predictors, could also be
7
8 targeted in the selective prevention programs and used in identifying or characterizing high-risk
9
10 participants. Although previous prevention programs have had limited success (Pratt &
11
12 Woolfenden, 2002), they would be improved by considering these gender- and development-
13
14 specific risk factors. We also recommend more studies that integrate gender and developmental
15
16 perspectives into research, which would be useful for elucidating the mechanisms of the
17
18 developmental psychopathology of disordered eating.
19
20
21

22 **Acknowledgments**

23
24 The authors wish to acknowledge and thank Stephanie Zerwas (assistant professor) and Cynthia
25
26 Bulik (professor) at the University of North Carolina for their valuable comments on drafts of
27
28 this paper.
29
30

31 **Funding**

32
33 This research was funded by the Research Council of Norway (grant # 196226V50).
34
35

36 **Declaration of Interest**

37
38 None.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- Alsaker, F. D., & Olweus, D. (1986). Assessment of global negative self-evaluations and perceived stability of self in Norwegian preadolescents and adolescents. *Journal of Early Adolescence, 6*, 269-278.
- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders* (4th ed., text revision). Washington, DC: American Psychiatric Press.
- Beato-Fernandez, L., Rodriguez-Cano, T., Belmonte-Llario, A., & Martinez-Delgado, C. (2004). Risk factors for eating disorders in adolescents: A Spanish community-based longitudinal study. *European Child & Adolescent Psychiatry, 13*, 287-294.
- Brown, T. A., Cash, T. F., & Lewis, R. J. (1989). Body-image disturbances in adolescent female binge purgers: A brief report of the results of a national survey in the U.S.A. *Journal of Child Psychology and Psychiatry and Allied Disciplines, 30*, 605-613.
- Campbell, J. D. (1990). Self-esteem and clarity of the self-concept. *Journal of Personality and Social Psychology, 59*, 538-549.
- Cicchetti, D., & Rogosch, F. A. (2002). A developmental psychopathology perspective on adolescence. *Journal of Consulting and Clinical Psychology, 70*, 6-20.
- Culbert, K. M., Racine, S. E., & Klump, K. L. (2011). The influence of gender and puberty on the heritability of disordered eating symptoms. In R. A. H. Adan & W. H. Kaye (Eds.), *Behavioral neurobiology of eating disorders: Vol. 6* (pp. 177-185). Heidelberg, Germany: Springer.
- Cummings, E. M., Davies, T. P., & Campbell, B. S. (2000). *Developmental psychopathology and family process: Theory, research, and clinical implications* (pp. 17-34). New York, NY: Guilford Press.

- 1
2
3
4 Dawson, J. F., & Richter, A. W. (2006). Probing three-way interactions in moderated multiple
5 regression: Development and application of a slope difference test. *Journal of applied*
6 *Psychology, 91*, 917-926.
7
8
9
10
11 De Goede, I. H. A., Branje, S. J. T., & Meeus, W. H. J. (2009). Developmental changes in
12 adolescents' perceptions of relationships with their parents. *Journal of Youth and*
13 *Adolescence, 38*, 75-88.
14
15
16
17
18 Derogatis, L. R., Lipman, R. S., Rickels, K., Uhlenhuth, E. H., & Covi, L. (1974). The Hopkins
19 Symptom Checklist (HSCL): A self-report symptom inventory. *Behavioral Sciences, 19*,
20 1-15.
21
22
23
24
25 Engelsen, B. K., & Hagtvet, K. A. (1999). The dimensionality of the 12-item version of the
26 Eating Attitudes Test: Confirmatory factor analyses. *Scandinavian Journal of*
27 *Psychology, 40*, 293-300.
28
29
30
31 Fairburn, C. G., & Harrison, P. J. (2003). Eating disorders. *The Lancet, 361*, 407-416.
32
33
34 Ferreiro, F., Seoane, G., & Senra, C. (2011). A prospective study of risk factors for the
35 development of depression and disordered eating in adolescents. *Journal of Clinical*
36 *Child & Adolescent Psychology, 40*, 500-505.
37
38
39
40 Fredrickson, B. L., & Roberts, T. A. (1997). Objectification theory: Toward understanding
41 women's lived experiences and mental health risks. *Psychology of Women Quarterly, 21*,
42 173-206.
43
44
45
46 Fredrickson, B. L., Roberts, T. A., Noll, S. M., Quinn, D. M., & Twenge, J. M. (1998). That
47 swimsuit becomes you: Sex differences in self-objectification, restrained eating, and math
48 performance. *Journal of Personality and Social Psychology, 75*, 269-284.
49
50
51
52
53 Garner, D. M., Olmsted, M. P., Bohr, Y., & Garfinkel, P. E. (1982). The eating attitudes test:
54 Psychometric features and clinical correlates. *Psychological Medicine, 12*, 871-878.
55
56
57
58
59
60

- 1
2
3
4 Gilligan, C. (1982). *In a different voice: Psychological theory and women's development* (pp. 39-
5
6
7 57). Cambridge, MA: Harvard University Press.
- 8
9 Goodman, E., Hinden, B. R., & Khandelwal, S. (2000). Accuracy of teen and parental reports of
10
11 obesity and body mass index. *Pediatrics, 106*, 52-58.
- 12
13 Graber, J. A., Brooks-Gunn, J., Paikoff, R. L., & Warren, M. P. (1994). Prediction of eating
14
15 problems: An 8-year study of adolescent girls. *Developmental Psychology, 30*, 823-834.
- 16
17
18 Haedt, A. A., & Keel, P. K. (2010). Comparing definitions of purging disorder on point
19
20 prevalence and associations with external validators. *International Journal of Eating*
21
22 *Disorders, 43*, 433-439.
- 23
24
25 Hankin, B. L., Abramson, L. Y., Moffitt, T. E., Silva, P. A., McGee, R., & Angell, K. E. (1998).
26
27 Development of depression from preadolescence to young adulthood: Emerging gender
28
29 differences in a 10-year longitudinal study. *Journal of Abnormal Psychology, 107*, 128-
30
31 140.
- 32
33 Harter, S. (1988). *Manual for the self-perception profile for adolescents*. Denver, CO: University
34
35 of Denver.
- 36
37
38 Hill, A. J. (1993). Pre-adolescent dieting: Implications for eating disorders. *International Review*
39
40 *of Psychiatry, 5*, 87-99.
- 41
42
43 Jacobi, C., Hayward, C., de Zwaan, M., Kraemer, H. C., & Agras, W. S. (2004). Coming to terms
44
45 with risk factors for eating disorders: Application of risk terminology and suggestions for
46
47 a general taxonomy. *Psychological Bulletin, 130*, 19-65.
- 48
49
50 Kansi, J., Wichstrom, L., & Bergman, L. R. (2003). Eating problems and the self-concept:
51
52 Results based on a representative sample of Norwegian adolescent girls. *Journal of*
53
54 *Youth and Adolescence, 32*, 325-335.
- 55
56
57 Keel, P. K., Fulkerson, J. A., & Leon, G. R. (1997). Disordered eating precursors in pre- and
58
59 early adolescent girls and boys. *Journal of Youth and Adolescence, 26*, 203-216.
- 60

- 1
2
3
4 Klump, K. L., Keel, P. K., Sisk, C., & Burt, S. A. (2010). Preliminary evidence that estradiol
5 moderates genetic influences on disordered eating attitudes and behaviors during puberty.
6
7 *Psychological Medicine, 40*, 1745-1753.
8
9
10
11 Kotler, L. A., Cohen, P., Davies, M., Pine, D. S., & Walsh, B. T. (2001). Longitudinal
12 relationships between childhood, adolescent, and adult eating disorders. *Journal of the*
13 *American Academy of Child & Adolescent Psychiatry, 40*, 1434-1440.
14
15
16
17 Lavik, N. J., Clausen, S. E., & Pedersen, W. (1991). Eating behaviour, drug use,
18 psychopathology and parental bonding in adolescents in Norway. *Acta Psychiatrica*
19 *Scandinavica, 84*, 387-390.
20
21
22
23
24 Le Grange, D., Lock, J., Loeb, K., & Nicholls, D. (2010). Academy for Eating Disorders Position
25 Paper: The role of the family in eating disorders. *International Journal of Eating*
26 *Disorders, 43*, 1-5.
27
28
29
30
31 Leon, G. R., Fulkerson, J. A., Perry, C. L., Keel, P. K., & Klump, K. L. (1999). Three to four
32 year prospective evaluation of personality and behavioral risk factors for later disordered
33 eating in adolescent girls and boys. *Journal of Youth and Adolescence, 28*, 181-196.
34
35
36
37
38 Levine, M. P. (2012). Loneliness and eating disorders. *Journal of Psychology, 146*, 243-257.
39
40 McCabe, M. P., & Ricciardelli, L. A. (2004). Body image dissatisfaction among males across the
41 lifespan: A review of past literature. *Journal of Psychosomatic Research, 56*, 675-685.
42
43
44 McCabe, M. P., & Ricciardelli, L. A. (2006). A prospective study of extreme weight change
45 behaviors among adolescent boys and girls. *Journal of Youth and Adolescence, 35*, 425-
46 434.
47
48
49
50
51 McCabe, M. P., & Vincent, M. A. (2003). The role of biodevelopmental and psychological
52 factors in disordered eating among adolescent males and females. *European Eating*
53 *Disorders Review, 11*, 315-328.
54
55
56
57
58
59
60

- 1
2
3
4 Mond, J., Hay, P., Rodgers, B., Owen, C., Crosby, R., & Mitchell, J. (2006). Use of extreme
5 weight control behaviours with and without binge eating in a community sample:
6
7 implications for the classification of bulimic-type eating disorders. *International Journal*
8
9 *of Eating Disorders*, 39, 294-302.
- 10
11
12 Murnen, S. K., & Smolak, L. (1997). Femininity, masculinity, and disordered eating: A meta-
13
14 analytic review. *International Journal of Eating Disorders*, 22, 231-242.
- 15
16
17 Murphy, E., Brewin, C. R., & Silka, L. (1997). The assessment of parenting using the parental
18
19 bonding instrument: Two or three factors? *Psychological Medicine*, 27, 333-341.
- 20
21
22 Neumark-Sztainer, D., Wall, M., Larson, N. I., Eisenberg, M. E., & Loth, K. (2011). Dieting and
23
24 disordered eating behaviors from adolescence to young adulthood: Findings from a 10-
25
26 year longitudinal study. *Journal of American Dietetic Association*, 111, 1004-1011.
- 27
28
29 Parker, G., Tupling, H., & Brown, L. B. (1979). A parental bonding instrument. *British Journal*
30
31 *of Medical Psychology*, 52, 1-10.
- 32
33
34 Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of
35
36 pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*,
37
38 17, 117-133.
- 39
40
41 Pratt, B. M., & Woolfenden, S. R. (2002). Interventions for preventing eating disorders in
42
43 children and adolescents. *Cochrane Database of Systematic Reviews*, Issue 2, Art. No.:
44
45 CD002891.
- 46
47
48 Rabe-Hesketh, S., & Skrondal, A. (2008). *Multilevel and longitudinal modeling using Stata* (2nd
49
50 ed.).(pp. 141-173). College Station, TX: Stata Press.
- 51
52
53 Ricciardelli, L. A., & McCabe, M. P. (2004). A biopsychosocial model of disordered eating and
54
55 the pursuit of muscularity in adolescent boys. *Psychological Bulletin*, 130, 179-205.
- 56
57
58
59
60

- 1
2
3
4 Rose, A. J., & Rudolph, K. D. (2006). A review of sex differences in peer relationship processes:
5
6 Potential trade-offs for the emotional and behavioral development of girls and boys.
7
8 *Psychological Bulletin*, 132, 98-131.
9
- 10
11 Rosen, K. (1996). The principles of developmental psychopathology: Illustration from the study
12
13 of eating disorders. In L. Smolak, M. P. Levine, & R. H. Striegel-Moore (Eds.), *The*
14
15 *developmental psychopathology of eating disorders: Implications for research,*
16
17 *prevention, and treatment* (pp. 3-30). Mahwah, NJ: Lawrence Erlbaum Associates.
18
19
- 20 Rosenberg, M. (1986). Self-concept from middle childhood through adolescence. In D. Suls & A.
21
22 G. Greenwald (Eds.), *Psychological perspectives on the self: Vol. 3* (pp. 107-136).
23
24 Hillsdale, NJ: Lawrence Erlbaum Associates.
25
- 26
27 Russell, D., Peplau, L. A., & Cutrona, C. E. (1980). The revised UCLA Loneliness Scale:
28
29 Concurrent and discriminant validity evidence. *Journal of Personality and Social*
30
31 *Psychology*, 39, 472-480.
32
- 33
34 Rutter, M. (1988). Epidemiological approaches to developmental psychopathology. *Archives of*
35
36 *General Psychiatry*, 45, 486-495.
37
- 38
39 Rutter, M. (1989). Age as an ambiguous variable in developmental research: Some
40
41 epidemiological considerations from developmental psychopathology. *International*
42
43 *Journal of Behavioral Development*, 12, 1-34.
44
- 45
46 Shisslak, C. M., Crago, M., & Estes, L. S. (1995). The spectrum of eating disturbances.
47
48 *International Journal of Eating Disorders*, 18, 209-219.
49
- 50
51 Shomaker, L. B., & Furman, W. (2009). Parent-adolescent relationship qualities, internal
52
53 working models, and attachment styles as predictors of adolescents' interactions with
54
55 friends. *Journal of Social and Personal Relationships*, 26, 579-603.
56
- 57
58 Simon, J., Schmidt, U., & Pilling, S. (2005). The health service use and cost of eating disorders.
59
60 *Psychological Medicine*, 35, 1543-1551.

- 1
2
3
4 Smink, F. R., van Hoeken, D., & Hoek, H. W. (2012). Epidemiology of eating disorders:
5
6 Incidence, prevalence and mortality rates. *Current Psychiatry Reports, 14*, 406-414.
7
8
9 Smolak, L., & Levine, M. P. (1996). Adolescent transitions and the development of eating
10
11 problems. In L. Smolak, M. P. Levine & R. H. Striegel-Moore (Eds.), *The developmental*
12
13 *psychopathology of eating disorders: Implications for research, prevention, and*
14
15 *treatment* (pp. 207-333). Hillsdale, NJ: Lawrence Erlbaum Associates.
16
17
18 Smolak, L., Levine, M. P., & Gralen, S. (1993). The impact of puberty and dating on eating
19
20 problems among middle school girls. *Journal of Youth and Adolescence, 22*, 355-368.
21
22
23 Smolak, L., & Striegel-Moore, R. (1996). The implications of developmental research for eating
24
25 disorders. In L. Smolak, M. P. Levine & R. H. Striegel-Moore (Eds.), *The developmental*
26
27 *psychopathology of eating disorders: Implications for research, prevention, and*
28
29 *treatment* (pp. 183-203). Hillsdale, NJ: Lawrence Erlbaum Associates.
30
31
32 Stice, E. (2002). Risk and maintenance factors for eating pathology: A meta-analytic review.
33
34 *Psychological Bulletin, 128*, 825-848.
35
36
37 Stice, E., Akutagawa, D., Gaggar, A., & Agras, W. S. (2000). Negative affect moderates the
38
39 relation between dieting and binge eating. *International Journal of Eating Disorders, 27*,
40
41 218-229.
42
43
44 Stice, E., Killen, J. D., Hayward, C., & Taylor, C. B. (1998). Support for the continuity
45
46 hypothesis of bulimic pathology. *Journal of Consulting and Clinical Psychology, 66*,
47
48 784-790.
49
50
51 Striegel-Moore, R. H., Fairburn, C. G., Wilfley, D. E., Pike, K. M., Dohm, F. A., & Kraemer, H.
52
53 C. (2005). Toward an understanding of risk factors for binge-eating disorder in black and
54
55 white women: A community-based case-control study. *Psychological Medicine, 35*, 907-
56
57 917.
58
59
60

- 1
2
3
4 Tata, P., Fox, J., & Cooper, J. (2001). An investigation into the influence of gender and parenting
5 styles on excessive exercise and disordered eating. *European Eating Disorders Review*, 9,
6 194-206.
7
8
9
10 Tomori, M., Zalar, B., & Plesnicar, B. K. (2000). Gender differences in psychosocial risk factors
11 among Slovenian adolescents. *Adolescence*, 35, 431-443.
12
13
14
15 Wichstrom, L. (1995). Harter's Self-Perception Profile for adolescents: Reliability, validity, and
16 evaluation of the question format. *Journal of Personality Assessment*, 65, 100-116.
17
18
19
20 Wilfley, D. E., Bishop, M. E., Wilson, G. T., & Agras, W. S. (2007). Classification of eating
21 disorders: Toward DSM-V. *International Journal of Eating Disorders*, 40 Suppl., S123-
22 S129.
23
24
25
26
27 Worell, J., & Todd, J. (1996). Development of the gendered self. In L. Smolak, M. P. Levine &
28 R. H. Striegel-Moore (Eds.), *The developmental psychopathology of eating disorders:
29 Implications for research, prevention, and treatment* (pp. 135-156). Hillsdale, NJ:
30 Lawrence Erlbaum Associates.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

Table 1

Summary Measures for Variables during Adolescence and Young Adulthood Phases, Males and Females Separately

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Variables	T1- Early adolescence		T1-Mid-adolescence		T2-Mid-adolescence		T2-Late adolescence		T3- Early young adulthood		T4-Middle young adulthood	
	Males N=1,268	Females N=1,291	Males N=1,448	Females N=1,672	Males N=1,268	Females N=1,291	Males N=1,448	Females N=1,672	Males N=1,217	Females N=1,528	Males N=1,201	Females N=1,517
	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)	M (SD)
Disordered eating	1.52(.36)	1.68(.39)	1.46(.30)	1.75(.37)	1.44(.33)	1.73(.38)	1.41(.31)	1.76(.39)	1.44(.29)	1.71(.38)	1.47(.35)	1.63(.39)
Age	13.49(.55)	13.45(.58)	16.11(.75)	16.12(.76)	15.09(.67)	15.04(.71)	17.76(.72)	17.75(.71)	21.83(1.71)	21.81(1.79)	28.36(1.67)	28.34(1.73)
Body mass index	19.30(2.57)	19.07(2.47)	21.05(2.54)	20.56(2.34)	20.98(5.82)	20.31(5.58)	22.21(4.35)	21.34(4.44)	24.14(4.78)	22.67(4.28)	25.29(3.75)	23.62(4.15)
Appearance satisfaction	3.61(.60)	3.24(.65)	3.68(.55)	3.23(.61)	3.71(.66)	3.20(.66)	3.75(.56)	3.27(.61)	3.71(.63)	3.32(.62)	3.64(.64)	3.38(.59)
Self-worth	2.97 (.55)	2.79(.57)	2.99(.54)	2.76(.53)	2.59(.35)	2.50(.33)	2.63(.32)	2.53(.33)	2.64(.13)	2.53(.30)	2.58(.30)	2.51(.28)
Negative affectivity	1.45(.41)	1.63(.48)	1.46(.37)	1.79(.52)	1.44(.41)	1.73(.52)	1.46(.41)	1.79(.52)	1.45(.43)	1.67(.52)	1.41(.43)	1.52(.48)
Loneliness	1.86(.53)	1.93(.58)	1.83(.53)	1.91(.54)	1.83(.57)	1.88(.58)	1.78(.54)	1.85(.54)	1.78(.53)	1.83(.48)	1.76(.51)	1.79(.49)
Alcohol intoxication	1.32(.82)	1.30(.78)	2.54(1.65)	2.53(1.56)	2.04(1.44)	2.05(1.43)	3.25(1.70)	3.05(1.56)	4.07(1.49)	3.53(1.48)	3.90(1.47)	3.12(1.51)
Illicit drug use	1.01(.21)	1.01(.15)	1.03(.34)	1.02(.30)	1.08(.43)	1.07(.41)	1.13(.49)	1.09(.38)	1.11(.56)	1.06(.41)	1.12(.56)	1.05(.37)
Self-concept instability	2.56(.63)	2.76(.59)	2.39(.65)	2.77(.61)	2.29(.71)	2.74(.63)	2.21(.70)	2.69(.68)	2.10(.73)	2.51(.73)	1.95(.70)	2.27(.72)
Pubertal development	2.21(.65)	2.78(.62)	0.76(1.33)	0.94(1.52)	-	-	-	-	-	-	-	-
Parental bonding styles:												
Over-protective	2.25(.52)	2.19(.54)	2.08(.54)	2.01(.58)	-	-	-	-	-	-	-	-
Care	3.12(.51)	3.19(.55)	3.10(.51)	3.15(.57)	-	-	-	-	-	-	-	-

Note. Pubertal development and parental bonding styles were measured at Time 1 only.

Table 2

Linear Regression Results for Early Predictors of Disordered Eating in Middle and Late Adolescence, Males and Females Separately

Predictors	Males				Females			
	Mid-adolescence N=1,268		Late adolescence N=1,448		Mid-adolescence N=1,291		Late adolescence N=1,672	
	β (SE)	$\beta+$ (SE)	β (SE)	$\beta+$ (SE)	β (SE)	$\beta+$ (SE)	β (SE)	$\beta+$ (SE)
Initial disordered eating	0.42(.04)***	0.37(.03)***	0.54(.04)***	0.45(.02)***	0.51(.01)***	0.46(.03)***	0.62(.03)***	0.60(.02)***
Body mass index (BMI)	0.11(.003)***	0.09(.003)**	0.09(.003)***	0.10(.003)***	0.09(.004)***	0.09(.03)**	0.04(.002)**	0.04(0.002)**
Appearance satisfaction	-0.14(.02)***	-0.02(.02)	-0.11(.01)***	-0.06(.01)**	-0.07(.02)*	0.02(.02)	-0.06(.01)**	-0.05(.01)*
Self-worth	-0.15(.02)***	-0.02(.02)	-0.11(.02)***	-0.02(0.02)	-0.10(.02)***	-0.08(.02)**	-0.06(.02)**	-0.003(.02)
Negative affectivity	0.15(.03)***	0.09(.02)**	0.12(.02)***	0.07(.02)**	0.04(.02)	-	0.05(.02)*	0.02(.02)
Loneliness	0.12(.02)***	0.05(.02)	0.13(.02)***	0.09(.01)**	0.02(.02)	-	0.001(.01)	-
Alcohol intoxication	0.004(.01)	-	0.04(.004)	-	-0.01(.01)	-	0.03(.005)	-
Illicit drug use	0.07(.04)**	0.06(.04)	0.01(.02)	-	-0.05(.06)*	-0.05(.06)	0.02(.03)	-
Self-concept instability	0.11(.02)***	0.05(.01)	0.06(.01)*	0.01(.01)	0.03(.02)	-	0.04(.01)**	0.05(.01)*
Pubertal development	0.04(.02)	-	0.08(.01)*	0.09(.01)**	0.03(.02)	-	0.03(.01)	-
Parental bonding styles:								
Over-protective	0.09(.02)***	-0.05(.02)	0.01(.01)	-	0.07(.02)**	0.06(.02)*	0.04(.01)*	0.03(.01)*
Care	-0.11(.02)***	0.03(.26)	-0.03(.01)	-	-0.02(.02)	-	-0.03(.01)	-


Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. β = standardized regression coefficients with standard errors in parentheses controlled for initial disordered eating and age. $\beta+$ = standardized regression coefficients with standard errors in parentheses from multiple linear regression models with control for other predictors, initial disordered eating, and age.

Table 3

Random Intercept Model Results for Predictors at T2 on Disordered Eating in Young Adulthood, Males and Females Separately

Predictors at T2	Males N=923		Females N=1,241	
	β (SE)	$\beta+$ (SE)	β (SE)	$\beta+$ (SE)
Initial disordered eating	0.36(.03)***	0.34(.03)***	0.38(.02)***	0.31(.02)***
Body mass index (BMI)	0.002(.002)	-	0.003(.002)	-
Appearance satisfaction	-0.03(.01)*	-0.01(.01)	-0.07(.01)***	-0.05(.01)***
Self-worth	-0.01(.02)	-	-0.05(.03)	-
Negative affectivity	0.11(.02)***	0.06(.02)**	0.06(.02)**	0.03(.02)
Loneliness	0.06(.01)***	0.02(.02)	0.04(.01)**	0.01(.01)
Alcohol intoxication	0.002(.005)	-	-0.004(.01)	-
Illicit drug use	-0.01(.02)	-	-0.02(.02)	-
Self-concept instability	0.05(.01)***	0.03(.01)**	0.05(.01)***	0.03(.01)**
Pubertal development	0.02(.01)	-	0.002(.01)	-
Parental bonding styles:				
Over-protective	0.002(.01)	-	0.01(.01)	-
Care	-0.04(.02)*	-0.01(.01)	0.001(.01)	-
Survey time points (T3 vs.T4)	0.03(.01)**	0.03(.01)**	-0.09(.01)***	-0.08(.01)***

Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. β = regression coefficients with standard errors in parenthesis controlled for age and disordered eating at T2. $\beta+$ = regression coefficients with standard errors in parentheses from multiple random intercept models with control for other predictors, age, and disordered eating at T2. Pubertal development and parental bonding styles were measured at T1 only.

Figure 1 Selected study population at each time point

212x87mm (96 x 96 DPI)