

On the normal sheaf of determinantal varieties

By *Jan O. Kleppe* at Oslo and *Rosa M. Miró-Roig* at Barcelona

Abstract. Let X be a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension c , i.e. a scheme defined by the maximal minors of a $t \times (t + c - 1)$ homogeneous polynomial matrix \mathcal{A} . In this paper, we study the main features of its normal sheaf \mathcal{N}_X . We prove that under some mild restrictions: (1) there exists a line bundle \mathcal{L} on $X \setminus \text{Sing}(X)$ such that $\mathcal{N}_X \otimes \mathcal{L}$ is arithmetically Cohen–Macaulay and, even more, it is Ulrich whenever the entries of \mathcal{A} are linear forms, (2) \mathcal{N}_X is simple (hence, indecomposable) and, finally, (3) \mathcal{N}_X is μ -(semi)stable provided the entries of \mathcal{A} are linear forms.

1. Introduction

The goal of this paper is to study the main properties of the normal sheaf of a determinantal scheme. Determinantal schemes have been a central topic in both Commutative Algebra and Algebraic Geometry. They are schemes defined by the vanishing of the $r \times r$ minors of a homogeneous polynomial matrix; and as a classical examples of determinantal schemes we have rational normal scrolls, Segre varieties and Veronese varieties. On the other hand, the normal sheaf \mathcal{N}_X of a projective scheme $X \subset \mathbb{P}^n$ has been intensively studied since it reflects many properties of the embedding. For instance, if $X \subset \mathbb{P}^n$ is a smooth projective variety of dimension $d \geq n/2$, $n \geq 5$, and \mathcal{N}_X splits into a sum of line bundles, then X is a complete intersection (cf. [2, Corollary 3.6]). This paper is entirely devoted to study the main features of the normal sheaf \mathcal{N}_X of a (linear) standard determinantal scheme $X \subset \mathbb{P}^n$, i.e. schemes defined by the maximal minors of a homogeneous polynomial matrix. In particular, we will study the Macaulayness, simplicity and μ -stability of \mathcal{N}_X .

Since the seminal work by Horrocks characterizing ACM sheaves on \mathbb{P}^n as those that completely split into a sum of line bundles (cf. [21]), a big amount of papers devoted to study ACM sheaves on projective schemes has appeared. Recall that a sheaf \mathcal{E} on a projective scheme $X \subset \mathbb{P}^n$ is arithmetically Cohen–Macaulay (ACM, for short) if it is locally Cohen–Macaulay and $H_*^i(\mathcal{E}) = \sum_{t \in \mathbb{Z}} H^i(X, \mathcal{E}(t)) = 0$ for $0 < i < \dim X$. It turns out that a natural way to measure the complexity of a projective scheme is to ask for the families of ACM sheaves that it supports. Mimicking an analogous trichotomy in Representation Theory, in [11] it was proposed a classification of ACM projective varieties as *finite*, *tame* or *wild* according to the

Second author partially supported by MTMM2010-15256.

complexity of their associated category of ACM vector bundles and it was proved that this trichotomy is exhaustive for the case of ACM curves: rational curves are finite, elliptic curves are tame and curves of higher genus are wild. Unfortunately very little is known for varieties of higher dimension.

Among ACM vector bundles \mathcal{E} on a given variety X , it is interesting to spot a very important subclass for which its associated module $\bigoplus_t H^0(X, \mathcal{E}(t))$ has the maximal number of generators, which turns out to be $\deg(X) \operatorname{rk}(\mathcal{E})$. The algebraic counterpart has also arisen a lot of interest. In fact, Ulrich proved that for a local ring R the minimal number $m(M)$ of generators of a maximal Cohen–Macaulay (MCM) module M is bounded by $m(M) \leq e(R) \operatorname{rk}(M)$ where $e(R)$ denotes the multiplicity of R (cf. [36]). MCM modules attaining this bound are called Ulrich modules. These algebraic considerations prompted to define, for a projective scheme $X \subset \mathbb{P}^n$, a sheaf \mathcal{E} on X to be Ulrich if it is ACM and the associated graded module $H_*^0(\mathcal{E})$ is Ulrich. When \mathcal{E} is initialized (i.e. $H^0(\mathcal{E}) \neq 0$ but $H^0(\mathcal{E}(-1)) = 0$), the last condition is equivalent to $\dim H^0(X, \mathcal{E}) = \deg(X) \operatorname{rk}(\mathcal{E})$. The search of Ulrich sheaves on a particular variety is a challenging problem. In fact, few examples of varieties supporting Ulrich sheaves are known, although in [15] it has been conjectured that any variety has an Ulrich sheaf. See [4, Proposition 2.8] for the existence of Ulrich bundles of rank one on linear standard determinantal schemes. Moreover, the recent interest in the existence of Ulrich sheaves relies among other things on the fact that a d -dimensional variety $X \subset \mathbb{P}^n$ supports an Ulrich sheaf (bundle) if and only if the cone of cohomology tables of coherent sheaves (resp. vector bundles) on X coincides with the cone of cohomology tables of coherent sheaves (resp. vector bundles) on \mathbb{P}^d ([14, Theorem 4.2]).

In this paper, we will address the following longstanding problems:

Problem 1.1. Let $X \subset \mathbb{P}^n$ be a (linear) standard determinantal scheme of codimension c and let \mathcal{N}_X be its normal sheaf:

- (1) Is there any invertible sheaf \mathcal{L} on $X \setminus \operatorname{Sing}(X)$ such that $\mathcal{N}_X \otimes \mathcal{L}$ is ACM (resp. Ulrich)? If so, how does the minimal free $\mathcal{O}_{\mathbb{P}^n}$ -resolution of $\mathcal{N}_X \otimes \mathcal{L}$ look like?
- (2) Is there any invertible sheaf \mathcal{L}' on $X \setminus \operatorname{Sing}(X)$ such that $\bigwedge^q \mathcal{N}_X \otimes \mathcal{L}'$ is ACM (resp. Ulrich)?
- (3) Is \mathcal{N}_X simple, or at least indecomposable?
- (4) Is \mathcal{N}_X μ -(semi)stable?

In our approach, we often use the well-known fact that a locally free sheaf on the smooth locus of a reduced normal Cohen–Macaulay variety X extends uniquely to a reflexive sheaf on X , or that the morphism (2.2) of this paper is an isomorphism under appropriate depth conditions. We will start our work analyzing whether the normal sheaf \mathcal{N}_X (and its exterior powers) to a standard determinantal scheme is ACM. It is well known that the normal sheaf \mathcal{N}_X to a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension c is ACM if $1 \leq c \leq 2$ but it is no longer true for $c \geq 3$. Our first goal will be to prove that under some weak restriction there exists an invertible sheaf \mathcal{L} on $X \setminus \operatorname{Sing}(X)$ (i.e. a coherent \mathcal{O}_X -module that is invertible on $X \setminus \operatorname{Sing}(X)$) such that $\mathcal{N}_X \otimes \mathcal{L}$ is ACM. Our next aim will be to prove that for linear standard determinantal schemes $X \subset \mathbb{P}^n$ satisfying some mild hypothesis, $\mathcal{N}_X \otimes \mathcal{L}$ is not only ACM but also Ulrich.

As Hartshorne and Casanellas pointed out in [9] there are few examples of indecomposable ACM bundles of arbitrarily high rank. So, once we know the Macaulayness of a suitable twist $\mathcal{N}_X \otimes \mathcal{L}$ of the normal sheaf \mathcal{N}_X to a standard determinantal scheme $X \subset \mathbb{P}^n$ we are led to ask if it is indecomposable. Concerning Problem 1.1 (3), we are able to prove that under some weak conditions \mathcal{N}_X is indecomposable and, even more, it is simple. As it is explained in Section 4, this last property works in a much more general set up (cf. Theorem 4.7). Another challenging problem is the existence of μ -(semi)stable ACM bundles of high rank on projective schemes since μ -(semi)stable ACM bundles of higher ranks are essentially unknown due to the lack of criteria to check it when the rank is high. In the last part of the paper, we analyze the μ -(semi)stability of the normal sheaf \mathcal{N}_X to a standard determinantal scheme X . We prove that it is μ -semistable provided the homogeneous matrix associated to X has linear entries and μ -stable if, in addition, we assume that X has codimension 2.

Next we outline the structure of the paper. Section 2 provides the background and basic results on (linear) standard determinantal schemes and the associated complexes needed in the sequel as well as the definition and main properties on ACM sheaves and Ulrich sheaves on a projective scheme. We refer to [6] and [13] for more details on standard determinantal schemes and to [10, 15, 31, 32] for more details on Ulrich sheaves. In Section 3, we address Problems 1.1 (1) and (2) and we prove our main results. Given a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension c associated to a homogeneous $t \times (t + c - 1)$ polynomial matrix \mathcal{A} , we denote by M the cokernel of the graded morphism defined by \mathcal{A} and we prove that, under some mild assumption, $\mathcal{N}_X \otimes \widetilde{S_{c-2}M}$ is ACM (cf. Theorem 3.6). As a by-product we obtain an $\mathcal{O}_{\mathbb{P}^n}$ -resolution of $\mathcal{N}_X \otimes \mathcal{L}$ which turns out to be pure and linear in the case of linear standard determinantal schemes (cf. Theorem 3.6 and Corollary 3.7). To prove it we use a generalization of the mapping cone process together with a careful analysis of possible cancelations of repeated summands in the mapping cone construction. Parts of the results of this section are inspired by ideas developed in [24] leading to isomorphisms

$$\mathrm{Ext}^1(\widetilde{M}, \widetilde{S_i M}) \cong \mathrm{Hom}(\mathcal{I}_X / \mathcal{I}_X^2, \widetilde{S_{i-1} M})$$

and to good estimates of the depth of $\mathrm{Ext}^1(M, S_i M)$. We also generalize these results to higher Ext-groups and to exterior powers of twisted normal sheaves (cf. Theorem 3.16). At the end of this section, we guess a result, analogous to Theorem 3.16 for the exterior power of the normal sheaf (cf. Conjectures 3.8 and 3.21) based on our computations with Macaulay2. Section 4 deals with Problem 1.1 (3) and we determine conditions under which the normal sheaf \mathcal{N}_X of a standard determinantal scheme $X \subset \mathbb{P}^n$ is simple and, hence, indecomposable (cf. Theorem 4.3). We also determine the cohomology and hence the depth of the conormal sheaf of X (cf. Proposition 4.1 and Corollaries 3.20 and 4.2). Finally, in Section 5, we face Problem 1.1 (4) and we prove that the normal sheaf \mathcal{N}_X of a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension c associated to a $t \times (t + c - 1)$ matrix with linear entries is always μ -semistable (cf. Theorem 5.3) and even more it is μ -stable when $c = 2$ and $n \geq 4$ (cf. Theorem 5.7).

Acknowledgement. The second author would like to thank A. Conca, L. Costa and J. Pons-Llopis for useful discussions on the subject. She also thanks Oslo and Akershus University College for its hospitality during her visit to Oslo in October 2011. The first author thanks the University of Barcelona for its hospitality during his visit to Barcelona in June 2012.

We also thank the referees for their comments and for pointing out that our first formulation of Lemma 3.1 was inaccurate.

Notation. Throughout this paper K will be an algebraically closed field of characteristic zero, $R = K[x_0, x_1, \dots, x_n]$, $\mathfrak{m} = (x_0, \dots, x_n)$ and $\mathbb{P}^n = \text{Proj}(R)$. Given a closed subscheme $X \subset \mathbb{P}^n$, we denote by \mathcal{I}_X its ideal sheaf, \mathcal{N}_X its normal sheaf and $I(X) = H_*^0(\mathbb{P}^n, \mathcal{I}_X)$ its saturated homogeneous ideal unless $X = \emptyset$, in which case we let $I(X) = \mathfrak{m}$. If X is equidimensional and Cohen–Macaulay of codimension c , we set $\omega_X = \text{Ext}_{\mathcal{O}_{\mathbb{P}^n}}^c(\mathcal{O}_X, \mathcal{O}_{\mathbb{P}^n})(-n-1)$ to be its canonical sheaf. Given a coherent sheaf \mathcal{E} on X , we denote the twisted sheaf $\mathcal{E} \otimes \mathcal{O}_X(l)$ by $\mathcal{E}(l)$. As usual, $H^i(X, \mathcal{E})$ stands for the cohomology groups, $h^i(X, \mathcal{E})$ for their dimension and

$$H_*^i(X, \mathcal{E}) = \bigoplus_{l \in \mathbb{Z}} H^i(X, \mathcal{E}(l)).$$

For any graded quotient A of R of codimension c , we let $I_A = \ker(R \twoheadrightarrow A)$ and we let $N_A = \text{Hom}_R(I_A, A)$ be the normal module. If A is Cohen–Macaulay of codimension c , we let $K_A = \text{Ext}_R^c(A, R)(-n-1)$ be its canonical module. When we write $X = \text{Proj}(A)$, we let $A = R_X := R/I(X)$ and $K_X = K_A$. If M is a finitely generated graded A -module, let $\text{depth}_J M$ denote the length of a maximal M -sequence in a homogeneous ideal J and let $\text{depth} M = \text{depth}_{\mathfrak{m}} M$.

2. Preliminaries

For convenience of the reader we include in this section the background and basic results on (linear) standard determinantal varieties as well as on arithmetically Cohen–Macaulay sheaves and Ulrich sheaves needed in the sequel.

2.1. Determinantal varieties. Let us start recalling the definition of arithmetically Cohen–Macaulay subschemes of \mathbb{P}^n .

Definition 2.1. A subscheme $X \subset \mathbb{P}^n$ is said to be arithmetically Cohen–Macaulay (briefly, ACM) if its homogeneous coordinate ring $R/I(X)$ is a Cohen–Macaulay ring, i.e. $\text{depth} R/I(X) = \dim R/I(X)$.

Thanks to the graded version of the Auslander–Buchsbaum formula (for any finitely generated R -module M), $\text{pd} M = n + 1 - \text{depth} M$, we deduce that a subscheme $X \subset \mathbb{P}^n$ is ACM if and only if $\text{pd} R/I(X) = \text{codim} X$. Hence, if $X \subset \mathbb{P}^n$ is a codimension c ACM subscheme, a graded minimal free R -resolution of $I(X)$ is of the form

$$0 \longrightarrow F_c \xrightarrow{\varphi_c} F_{c-1} \xrightarrow{\varphi_{c-1}} \dots \xrightarrow{\varphi_2} F_1 \xrightarrow{\varphi_1} I(X) \longrightarrow 0$$

where

$$F_i = \bigoplus_{j \in \mathbb{Z}} R(-j)^{\beta_{i,j}}, \quad i = 1, \dots, c$$

(in this setting, minimal means that $\text{im}(\varphi_i) \subset \mathfrak{m} F_{i-1}$).

We now collect the results on standard determinantal schemes and the associated complexes needed in the sequel and we refer to [6] and [13] for more details.

Definition 2.2. If \mathcal{A} is a homogeneous matrix, we denote by $I(\mathcal{A})$ the ideal of R generated by the maximal minors of A and by $I_j(\mathcal{A})$ the ideal generated by the $j \times j$ minors of \mathcal{A} . A codimension c subscheme $X \subset \mathbb{P}^n$ is called a *standard determinantal scheme* if $I(X) = I(\mathcal{A})$ for some $t \times (t + c - 1)$ homogeneous matrix \mathcal{A} . In addition, we will say that X is a *linear standard determinantal scheme* if all entries of \mathcal{A} are linear forms.

As examples of standard determinantal schemes we have any ACM variety $X \subset \mathbb{P}^n$ of codimension 2 and as examples of linear determinantal schemes we have, for instance, rational normal curves, Segre varieties and rational normal scrolls. Now we are going to describe some complexes associated to a codimension c standard determinantal scheme X . To this end, we denote by $\varphi : F \rightarrow G$ the morphism of free graded R -modules of rank $t + c - 1$ and t , defined by the homogeneous matrix \mathcal{A} of X . We denote by $\mathcal{C}_i(\varphi)$ the (generalized) Koszul complex

$$\mathcal{C}_i(\varphi) : 0 \rightarrow \bigwedge^i F \otimes S_0(G) \rightarrow \bigwedge^{i-1} F \otimes S_1(G) \rightarrow \cdots \rightarrow \bigwedge^0 F \otimes S_i(G) \rightarrow 0$$

where $\bigwedge^i F$ stands for the i -th exterior power of F and $S_i(G)$ denotes the i -th symmetric power of G . Let $\mathcal{C}_i(\varphi)^*$ be the R -dual of $\mathcal{C}_i(\varphi)$. The dual map φ^* induces graded morphisms

$$\mu_i : \bigwedge^{t+i} F \otimes \bigwedge^t G^* \rightarrow \bigwedge^i F.$$

They can be used to splice the complexes $\mathcal{C}_{c-i-1}(\varphi)^* \otimes \bigwedge^{t+c-1} F \otimes \bigwedge^t G^*$ and $\mathcal{C}_i(\varphi)$ to a complex $\mathcal{D}_i(\varphi)$:

$$\begin{aligned} (2.1) \quad 0 &\rightarrow \bigwedge^{t+c-1} F \otimes S_{c-i-1}(G)^* \otimes \bigwedge^t G^* \\ &\rightarrow \bigwedge^{t+c-2} F \otimes S_{c-i-2}(G)^* \otimes \bigwedge^t G^* \\ &\rightarrow \cdots \rightarrow \bigwedge^{t+i} F \otimes S_0(G)^* \otimes \bigwedge^t G^* \\ &\xrightarrow{\epsilon_i} \bigwedge^i F \otimes S_0(G) \rightarrow \bigwedge^{i-1} F \otimes S_1(G) \rightarrow \cdots \rightarrow \bigwedge^0 F \otimes S_i(G) \rightarrow 0. \end{aligned}$$

The complex $\mathcal{D}_0(\varphi)$ is called the Eagon–Northcott complex and the complex $\mathcal{D}_1(\varphi)$ is called the Buchsbaum–Rim complex. Let us rename the complex $\mathcal{C}_c(\varphi)$ as $\mathcal{D}_c(\varphi)$. Denote by $I_m(\varphi)$ the ideal generated by the $m \times m$ minors of the matrix \mathcal{A} representing φ . Then, we have the following well-known result:

Proposition 2.3. *Let $X \subset \mathbb{P}^n$ be a standard determinantal subscheme of codimension c associated to a graded minimal (i.e. $\text{im}(\varphi) \subset \mathfrak{m}G$) morphism $\varphi : F \rightarrow G$ of free R -modules of rank $t + c - 1$ and t , respectively. Set $M = \text{Coker}(\varphi)$. Then, it holds:*

- (i) $\mathcal{D}_i(\varphi)$ is acyclic for $-1 \leq i \leq c$.
- (ii) $\mathcal{D}_0(\varphi)$ is a minimal free graded R -resolution of $R/I(X)$ and $\mathcal{D}_i(\varphi)$ is a minimal free graded R -resolution of length c of $S_i(M)$, $1 \leq i \leq c$.
- (iii) Up to twist, $K_X \cong S_{c-1}(M)$. So, up to twist, $\mathcal{D}_{c-1}(\varphi)$ is a minimal free graded R -module resolution of K_X .
- (iv) $\mathcal{D}_i(\varphi)$ is a minimal free graded R -resolution of $S_i(M)$ for $c + 1 \leq i$ whenever one has $\text{depth}_{I_m(\varphi)} R \geq t + c - m$ for every m such that $t \geq m \geq \max(1, t + c - i)$.

Proof. See for instance [6, Theorem 2.20] and [13, Theorem A2.10, Corollaries A2.12 and A2.13]. \square

It immediately follows from Proposition 2.3 (ii) that standard determinantal schemes are ACM. Let us also recall the following useful comparison of cohomology groups. If $Z \subset X$ is a closed subset such that $U = X \setminus Z$ is a local complete intersection, L, N are finitely generated $R/I(X)$ -modules, \tilde{N} is locally free on U and $\text{depth}_{I(Z)} L \geq r + 1$, then the natural map

$$(2.2) \quad \text{Ext}_{R/I(X)}^i(N, L) \rightarrow H_*^i(U, \mathcal{H}om_{\mathcal{O}_X}(\tilde{N}, \tilde{L}))$$

is an isomorphism, (resp. an injection) for $i < r$ (resp. $i = r$) cf. [20, Exposé VI]. Recall that we interpret $I(Z)$ as \mathfrak{m} if $Z = \emptyset$.

We end this subsection describing the Picard group of a smooth standard determinantal scheme X . Assume that $X \subset \mathbb{P}^n$ is given by the maximal minors of a $t \times (t + c - 1)$ homogeneous matrix \mathcal{A} representing a homomorphism φ of free graded R -modules

$$\varphi : F = \bigoplus_{i=1}^{t+c-1} R(-a_i) \rightarrow G = \bigoplus_{j=1}^t R(-b_j).$$

Without loss of generality, we may assume $a_1 \leq a_2 \leq \dots \leq a_{t+c-1}$ and $b_1 \leq b_2 \leq \dots \leq b_t$. Denote by H the general hyperplane section of X and by $Z \subset X$ the codimension 1 subscheme of X defined by the maximal minors of the $(t - 1) \times (t + c - 1)$ matrix obtained deleting the last row of \mathcal{A} . The following theorem computes the Picard group of X . Indeed, we have:

Theorem 2.4. *Let $X \subset \mathbb{P}^n$ be a smooth standard determinantal scheme of codimension $c \geq 2$. Set*

$$\ell := \sum_{j=1}^{t+c-1} a_j - \sum_{i=1}^t b_i.$$

Assume $t > 1$. If $n - c > 2$ and $a_1 - b_t > 0$, or if $n - c = 2$, $a_1 - b_t > 0$ and $\ell \geq n + 1$, then

$$\text{Pic}(X) \cong \mathbb{Z}^2 \cong \langle H, Z \rangle.$$

Proof. See [12, Corollary 2.4] for smooth standard determinantal varieties $X \subset \mathbb{P}^n$ of dimension $d \geq 3$ and [18, Proposition 5.2] for the case $d = 2$ (see also [30, Theorem II.4.2] for smooth surfaces $X \subset \mathbb{P}^4$). \square

2.2. ACM and Ulrich sheaves. We set up here some preliminary notions mainly concerning the definitions and basic results on ACM and Ulrich sheaves needed later.

Definition 2.5. Let $X \subset \mathbb{P}^n$ be a projective scheme and let \mathcal{E} be a coherent sheaf on X . Then \mathcal{E} is said to be *Arithmetically Cohen–Macaulay* (shortly, ACM) if it is locally Cohen–Macaulay (i.e., $\text{depth } \mathcal{E}_x = \dim \mathcal{O}_{X,x}$ for every point $x \in X$) and has no intermediate cohomology, i.e.

$$H^i(X, \mathcal{E}(t)) = 0 \quad \text{for all } t \text{ and } i = 1, \dots, \dim X - 1.$$

Notice that when X is a non-singular variety, any coherent ACM sheaf on X is locally free. A seminal result due to Horrocks (cf. [21]) asserts that, up to twist, there is only one indecomposable ACM bundle on \mathbb{P}^n , namely $\mathcal{O}_{\mathbb{P}^n}$. Ever since this result was stated, the study

of the category of indecomposable arithmetically Cohen–Macaulay bundles on a given projective scheme X has raised a lot of interest since it is a natural way to understand the complexity of the underlying variety X (for more information the reader can see [9, 10, 31–33]). One of the goals of this paper is to study whether the normal sheaf \mathcal{N}_X and the exterior power $\bigwedge^q \mathcal{N}_X$ of the normal sheaf (or suitable twists $\mathcal{N}_X \otimes \mathcal{L}$ and $\bigwedge^q \mathcal{N}_X \otimes \mathcal{L}'$ by invertible sheaves \mathcal{L} and \mathcal{L}') of a standard determinantal scheme $X \subset \mathbb{P}^n$ are ACM. ACM sheaves are closely related to their algebraic counterpart, the maximal Cohen–Macaulay modules.

Definition 2.6. A graded R_X -module E is a *maximal Cohen–Macaulay module* (MCM for short) if $\text{depth } E = \dim E = \dim R_X$.

In fact, we have:

Proposition 2.7. *Let $X \subseteq \mathbb{P}^n$ be an ACM scheme. There exists a bijection between ACM sheaves \mathcal{E} on X and MCM R_X -modules E given by the functors $E \rightarrow \widetilde{E}$ and $\mathcal{E} \rightarrow H_*^0(X, \mathcal{E})$.*

Proof. See [8, Proposition 2.1]. □

Definition 2.8. Given a closed subscheme $X \subset \mathbb{P}^n$, a coherent sheaf \mathcal{E} on X is said to be *initialized* if

$$H^0(X, \mathcal{E}(-1)) = 0 \quad \text{but} \quad H^0(X, \mathcal{E}) \neq 0.$$

Notice that when \mathcal{E} is a locally Cohen–Macaulay sheaf, there always exists an integer k such that $\mathcal{E}_{\text{init}} := \mathcal{E}(k)$ is initialized.

Let us now introduce the notion of Ulrich sheaf.

Definition 2.9. Given a projective scheme $X \subset \mathbb{P}^n$ and a coherent sheaf \mathcal{E} on X , we say that \mathcal{E} is an *Ulrich sheaf* if \mathcal{E} is an ACM sheaf and $h^0(\mathcal{E}_{\text{init}}) = \deg(X) \text{rk}(\mathcal{E})$.

We have the following result that justifies this definition:

Theorem 2.10. *Let $X \subseteq \mathbb{P}^n$ be an integral subscheme and let \mathcal{E} be an ACM sheaf on X . Then the minimal number of generators $m(\mathcal{E})$ of the R_X -module $H_*^0(\mathcal{E})$ is bounded by*

$$m(\mathcal{E}) \leq \deg(X) \text{rk}(\mathcal{E}).$$

Therefore, since it is obvious that for an initialized sheaf \mathcal{E} , $h^0(\mathcal{E}) \leq m(\mathcal{E})$, the minimal number of generators of Ulrich sheaves is as large as possible. Modules attaining this upper bound were studied by Ulrich in [36]. A detailed account is provided in [15]. In particular we have:

Theorem 2.11. *Let $X \subseteq \mathbb{P}^N$ be an n -dimensional ACM variety and let \mathcal{E} be an initialized ACM coherent sheaf on X . The following conditions are equivalent:*

- (i) \mathcal{E} is Ulrich.
- (ii) \mathcal{E} admits a linear $\mathcal{O}_{\mathbb{P}^N}$ -resolution of the form:

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^N}(-N+n)^{a_{N-n}} \rightarrow \cdots \rightarrow \mathcal{O}_{\mathbb{P}^N}(-1)^{a_1} \rightarrow \mathcal{O}_{\mathbb{P}^N}^{a_0} \rightarrow \mathcal{E} \rightarrow 0.$$

- (iii) $H^i(\mathcal{E}(-i)) = 0$ for $i > 0$ and $H^i(\mathcal{E}(-i-1)) = 0$ for $i < n$.
- (iv) For some (resp. all) finite linear projections $\pi : X \rightarrow \mathbb{P}^n$, the sheaf $\pi_*\mathcal{E}$ is the trivial sheaf $\mathcal{O}_{\mathbb{P}^n}^t$ for some t .

In particular, initialized Ulrich sheaves are 0-regular and therefore they are globally generated.

Proof. See [15, Proposition 2.1]. □

In the next section, we will prove that under some weak restrictions the normal sheaf \mathcal{N}_X of a linear standard determinantal scheme X twisted by a suitable invertible sheaf \mathcal{L} on $X \setminus \text{Sing}(X)$ is Ulrich (cf. Theorem 3.6); and as an immediate consequence we will deduce the μ -semistability of the normal sheaf of a linear standard determinantal scheme (cf. Theorem 5.3).

3. The Cohen–Macaulayness of the normal sheaf of a determinantal variety

The main goal of this section is to answer Problem 1.1 (1). More precisely, we prove that, under some mild assumptions, given a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension c there always exists an invertible sheaf \mathcal{L} on $X \setminus \text{Sing}(X)$ such $\mathcal{N}_X \otimes \mathcal{L}$ is ACM (cf. Theorem 3.6) and as a by-product we obtain an $\mathcal{O}_{\mathbb{P}^n}$ -resolution of $\mathcal{N}_X \otimes \mathcal{L}$ which turns out to be pure and linear in the case of linear standard determinantal schemes (cf. Theorem 3.6 and Corollary 3.7). At the end of this section, we conjecture an analogous result for the exterior power of the normal sheaf (cf. Conjecture 3.21).

In this section we use the following notation:

- $X \subset \mathbb{P}^n$ is a standard determinantal scheme of codimension c ,
- \mathcal{A} is the $t \times (t + c - 1)$ homogeneous matrix associated to X ,
- $I = I_t(\mathcal{A})$, $A = R/I$,
- $N_A := \text{Hom}_A(I/I^2, A)$, $\mathcal{N}_X := \widetilde{N}_A$,
- $\varphi : F := \bigoplus_{j=1}^{t+c-1} R(-a_j) \rightarrow G := \bigoplus_{i=1}^t R(-b_i)$ is the morphism of free R -modules associated to \mathcal{A} and $M := \text{coker}(\varphi)$.

We will assume $t > 1$, since the case $t = 1$ corresponds to a codimension c complete intersection $X \subset \mathbb{P}^n$ and its normal sheaf is well understood ($\mathcal{N}_X \cong \bigoplus_{i=1}^c \mathcal{O}_X(d_i)$). If $t \geq 2$, then \widetilde{M} is a locally free \mathcal{O}_X -module of rank 1 over $T := X \setminus V(J)$ where $J := I_{t-1}(\mathcal{A})$ and $T \hookrightarrow \mathbb{P}^n$ is a local complete intersection. Recall also that if $a_j > b_i$ for any i, j , then one has $V(J) = \text{Sing}(X)$, $\text{codim}_X(\text{Sing}(X)) = c + 2$ or $\text{Sing}(X) = \emptyset$ and $\text{codim}_{\mathbb{P}^n} X = c$ for a general choice of $\varphi \in \text{Hom}(F, G)$.

Recall that the normal sheaf \mathcal{N}_X of a standard determinantal scheme is ACM if and only if $H_*^i(\mathcal{N}_X) = 0$ for $0 < i < \dim X$. In particular, the normal sheaf \mathcal{N}_X of a standard determinantal curve $X \subset \mathbb{P}^n$ ($c = n - 1$) is always ACM as well as the normal sheaf \mathcal{N}_X of a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension $1 \leq c \leq 2$. Indeed, if $c = 1$, then $\mathcal{N}_X \cong \mathcal{O}_X(\delta)$ where $\delta := \deg(X)$ and if $c = 2$, then there is an exact sequence (cf. [28, equation (26)])

$$(3.1) \quad 0 \rightarrow \widetilde{F} \otimes_R \widetilde{G}^* \rightarrow ((\widetilde{F}^* \otimes_R \widetilde{F}) \oplus (\widetilde{G}^* \otimes_R \widetilde{G}))/R \rightarrow \widetilde{G} \otimes_R \widetilde{F}^* \rightarrow \mathcal{N}_X \rightarrow 0.$$

Unfortunately, it is no longer true for higher codimension and we only have that under some mild conditions $H_*^i(\mathcal{N}_X) = 0$ for $1 \leq i \leq n - c - 2$ (see [29, Lemma 35] for $c = 3$, [25, Corollary 5.5] for $3 \leq c \leq 4$ and [24, Theorem 5.11] for the general case). In this section we are going to prove that under some weak restrictions $\mathcal{N}_X \otimes \widehat{S}_{c-2}M$ is ACM, i.e.

$$H_*^i(X, \mathcal{N}_X \otimes \widehat{S}_{c-2}M) = 0 \quad \text{for } 1 \leq i \leq n - c - 1.$$

To prove it, we will use the following technical lemma which can be seen as a generalization of the mapping cone process and we include a proof for the sake of completeness. In this lemma the differentials of a complex, say Q_\bullet , are denoted by $d_Q^i : Q_i \rightarrow Q_{i-1}$.

Lemma 3.1. *Let $Q_\bullet \xrightarrow{\sigma_\bullet} P_\bullet \xrightarrow{\tau_\bullet} F_\bullet$ be morphisms of complexes satisfying*

$$Q_j = P_j = F_j = 0 \quad \text{for } j < 0$$

and assume that all three complexes are acyclic (exact for $j \neq 0$) and that the sequence

$$0 \rightarrow \text{coker } d_Q^1 \rightarrow \text{coker } d_P^1 \xrightarrow{\alpha} \text{coker } d_F^1$$

is exact. Moreover assume that there exists a morphism $\ell_\bullet : Q_\bullet \rightarrow F_\bullet[1]$ such that for any integer i ,

$$(3.2) \quad d_F^{i+1} \ell_i + \ell_{i-1} d_Q^i = \tau_i \sigma_i.$$

Then, the complex $Q_\bullet \oplus P_\bullet[1] \oplus F_\bullet[2]$ given by

$$Q_i \oplus P_{i+1} \oplus F_{i+2} \xrightarrow{d_{Q,P,F}^i} Q_{i-1} \oplus P_i \oplus F_{i+1}$$

where

$$d_{Q,P,F}^i := \begin{pmatrix} d_Q^i & 0 & 0 \\ \sigma_i & -d_P^{i+1} & 0 \\ \ell_i & -\tau_{i+1} & d_F^{i+2} \end{pmatrix}$$

is acyclic (exact for $i \neq -2$) and $\text{coker } d_{Q,P,F}^{-1} = \text{coker } \alpha$.

Proof. It is straightforward to show that $d_{Q,P,F}^{i-1} \circ d_{Q,P,F}^i = 0$ by using (3.2) and that the differentials of the complexes commute with σ_\bullet and τ_\bullet . To see the acyclicity of the complex $Q_\bullet \oplus P_\bullet[1] \oplus F_\bullet[2]$, let

$$\begin{pmatrix} d_Q^{i-1} & 0 & 0 \\ \sigma_{i-1} & -d_P^i & 0 \\ \ell_{i-1} & -\tau_i & d_F^{i+1} \end{pmatrix} \begin{pmatrix} q \\ p \\ f \end{pmatrix} = \begin{pmatrix} d_Q^{i-1}(q) \\ \sigma_{i-1}(q) - d_P^i(p) \\ \ell_{i-1}(q) - \tau_i(p) + d_F^{i+1}(f) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

We must show the existence of an element $(q', p', f') \in Q_i \oplus P_{i+1} \oplus F_{i+2}$ whose transpose maps to $(q, p, f)^\text{tr}$ via $d_{Q,P,F}^i$. Since Q_\bullet is exact for $i \neq 0$, there exists an element $q' \in Q_i$ such that $d_Q^i(q') = q$ provided $i > 1$. For $i = 1$ we use $\sigma_0(q) = d_P^1(p)$ and the injectivity of $\text{coker } d_Q^1 \rightarrow \text{coker } d_P^1$ to see that q maps to zero in $\text{coker } d_Q^1$. Hence there exists an element $q' \in Q_1$ such that $d_Q^1(q') = q$. We will treat the case $i = 0$ shortly.

Suppose $i > 0$. Since we have $\sigma_{i-1}(q) - d_P^i(p) = 0$, $q = d_Q^i(q')$ and $\sigma_{i-1}d_Q^i = d_P^i\sigma_i$, we get

$$d_P^i(\sigma_i(q') - p) = 0.$$

By the exactness of P_\bullet , there exists an element $p' \in P_{i+1}$ such that

$$d_P^{i+1}(p') = \sigma_i(q') - p.$$

Then we insert $p = \sigma_i(q') - d_P^{i+1}(p')$ and $q = d_Q^i(q')$ into $\ell_{i-1}(q) - \tau_i(p) + d_F^{i+1}(f) = 0$ and we use equation (3.2) and that d_P^{i+1} and d_F^{i+1} commute with τ_\bullet to conclude that

$$d_F^{i+1}(-\ell_i(q') + \tau_{i+1}(p') + f) = 0.$$

Hence there exists an element $f' \in F_{i+2}$ such that

$$f = \ell_i(q') - \tau_{i+1}(p') + d_F^{i+2}(f').$$

The expressions for q , p and f above prove precisely what we needed to show if $i > 0$.

Now suppose $i = 0$. Since $d_Q^{-1} = d_Q^0 = d_P^0 = \sigma_{-1} = \ell_{-1} = 0$ and $q \in Q_{-1} = 0$, we have an element $(p, f) \in P_0 \oplus F_1$ that maps to $-\tau_0(p) + d_F^1(f) = 0$ in F_0 . Since $\tau_0(p)$ maps to zero in $\text{coker } d_F^1$, it follows that p is sent to an element in $\text{coker } d_P^1$ that is contained in $\text{coker } d_Q^1$ by the exactness assumption on the sequence of cokernels of Lemma 3.1. Hence there exists an element $q' \in Q_0$ such that $\sigma_0(q') - p$ maps to zero in $\text{coker } d_P^1$, whence there is an element $p' \in P_1$ such that

$$d_P^1(p') = \sigma_0(q') - p.$$

Since we have $d_F^1(-\ell_0(q') + \tau_1(p') + f) = 0$, there exists an element $f' \in F_2$ such that

$$d_F^2(f') = -\ell_0(q') + \tau_1(p') + f,$$

and we get the expressions for p and f which we wanted to show.

Finally using that $(p, f) \in P_0 \oplus F_1$ maps to $-\tau_0(p) + d_F^1(f)$ in F_0 we easily get the identity $\text{coker } d_{Q,P,F}^{-1} = \text{coker } \alpha$ and we are done. \square

Theorem 3.2. *We keep the notation introduced above and, in addition, we assume that $\text{depth}_j A \geq 2$. Then:*

- $\text{Ext}_R^1(M, S_{c-1}M)$ is a maximal Cohen–Macaulay A -module of rank c .
- If $a_j = 1$ for all j and $b_i = 0$ for all i , then $\text{Ext}_R^1(M, S_{c-1}M)$ is an Ulrich sheaf of rank c .

Proof. Our primary goal is to show that $\text{Ext}_R^1(M, S_{c-1}M)$ is an MCM A -module by using the exact sequence

$$(3.3) \quad 0 \rightarrow S_{c-2}M \rightarrow G^* \otimes S_{c-1}M \rightarrow F^* \otimes S_{c-1}M \rightarrow \text{Ext}_R^1(M, S_{c-1}M) \rightarrow 0.$$

and Lemma 3.1 to exhibit a minimal free resolution of $\text{Ext}_R^1(M, S_{c-1}M)$ having length c . So let us start proving the existence of (3.3). Indeed, we look at the Buchsbaum–Rim complex (see Proposition 2.3 (ii))

$$(3.4) \quad \cdots \longrightarrow \bigwedge^{t+1} F \otimes \bigwedge^t G^* \otimes S_0 G^* \xrightarrow{\epsilon_1^*} F \xrightarrow{\varphi} G \longrightarrow M \longrightarrow 0.$$

Since $I_t(\varphi) \cdot M = 0$ and $\text{im}(\epsilon_1^*) \subseteq I_t(\varphi) \cdot F$, we get that the induced map

$$\text{Hom}_R(\epsilon_1^*, S_{c-1}M) = 0.$$

Therefore, the exact sequence (3.3) comes from applying the functor $\text{Hom}_R(-, S_{c-1}M)$ to the exact sequence (3.4), because under the assumption $\text{depth}_J A \geq 2$, we have $\text{depth}_J S_i M \geq 2$ for any $1 \leq i \leq c$ and hence

$$(3.5) \quad \begin{aligned} \text{Hom}_R(M, S_i M) &\cong H_*^0(X \setminus V(J), \mathcal{H}om_{\mathcal{O}_X}(\widetilde{M}, \widetilde{M}^{\otimes(i)})) \\ &\cong H_*^0(X \setminus V(J), \widetilde{M}^{\otimes(i-1)}) \\ &\cong S_{i-1}M. \end{aligned}$$

From the exact sequence (3.3) we deduce that $\text{Ext}_R^1(M, S_{c-1}M)$ has rank c ; let us prove that it is a maximal Cohen–Macaulay A -module. The idea will be to apply Lemma 3.1 to the following diagram which we will define as an expansion of (3.3) (we set $\bigwedge^i := \bigwedge^i F$).




Diagram A

Let us call Q_\bullet , P_\bullet and F_\bullet the resolutions of $S_{c-2}M$, $G^* \otimes S_{c-1}M$ and $F^* \otimes S_{c-1}M$, respectively. We need to define morphisms of complexes: $\sigma_\bullet : Q_\bullet \rightarrow P_\bullet$ and $\ell_\bullet : Q_\bullet \rightarrow F_\bullet[1]$ satisfying all the hypothesis of Lemma 3.1. Let us first recall the definition of

$$\partial_q^p : S_p G \otimes \bigwedge^q F \rightarrow S_{p+1} G \otimes \bigwedge^{q-1} F.$$

To this end, we take $\{x_i\}_{i=1}^t$ an R -free basis of G and let $\{x_i^*\}_{i=1}^t$ be the dual basis of G^* . According to [13, p. 592], the map ∂_q^p takes an element

$$m \otimes f \in S_p G \otimes \bigwedge^q F$$

to the element

$$\sum_{i=1}^t x_i m \otimes \varphi^*(x_i^*)(f) \in S_{p+1} G \otimes \bigwedge^{q-1} F.$$

For any integer i , $0 \leq i \leq c-2$, we define

$$\sigma_i : S_{c-2-i} G \otimes \bigwedge^i F \rightarrow G^* \otimes S_{c-1-i} G \otimes \bigwedge^i F$$

sending an element

$$m \otimes f \in S_{c-2-i} G \otimes \bigwedge^i F$$

to

$$\sigma_i(m \otimes f) = \sum_{j=1}^t x_j^* \otimes x_j m \otimes f \in G^* \otimes S_{c-1-i} G \otimes \bigwedge^i F.$$

It is easy to check that the following diagram commutes for any integer i , $0 \leq i \leq c-3$:

$$\begin{array}{ccc} S_{c-3-i} G \otimes \bigwedge^{i+1} F & \xrightarrow{\sigma_{i+1}} & G^* \otimes S_{c-2-i} G \otimes \bigwedge^{i+1} F \\ \downarrow \partial_{i+1}^{c-3-i} & & \downarrow 1_{G^*} \otimes \partial_{i+1}^{c-2-i} \\ S_{c-2-i} G \otimes \bigwedge^i F & \xrightarrow{\sigma_i} & G^* \otimes S_{c-1-i} G \otimes \bigwedge^i F. \end{array}$$

In this setting, we point out that the definition of σ_{c-2} implies the commutativity of the diagram

$$\begin{array}{ccc} S_0 G \otimes \bigwedge^{c-2} F & \xrightarrow{\sigma_{c-2}} & G^* \otimes S_1 G \otimes \bigwedge^{c-2} F \\ \downarrow \cong & & \downarrow \cong \\ R \otimes \bigwedge^{c-2} F & \xrightarrow{\text{tr} \otimes 1} & G^* \otimes G \otimes \bigwedge^{c-2} F, \end{array}$$

i.e., the map σ_{c-2} is induced by the trace map $R \rightarrow G^* \otimes G$ that is dual to the evaluation map $G^* \otimes G \rightarrow R$ (i.e. $\sigma_{c-2} = \text{tr} \otimes 1$). We will now define σ_c and σ_{c-1} in such a way that the two left upper squares of Diagram A commute or anticommute. Dualizing (i.e. applying $\text{Hom}_R(-, R)$) and using the isomorphism

$$\bigwedge^i F^* \cong \bigwedge^{t+c-1-i} F \otimes \bigwedge^{t+c-1} F^* \cong \bigwedge^{t+c-1-i} F,$$

it will be sufficient to prove the commutativity of the following diagram:

$$\begin{array}{ccc}
 G & \xleftarrow{\sigma_c^*} & G \otimes R \\
 \uparrow \varphi & & \uparrow \epsilon_0^* \\
 F & \xleftarrow{\sigma_{c-1}^*} & G \otimes S_0 G^* \otimes \bigwedge^t F \otimes \bigwedge^t G^* \\
 \uparrow (-1)^{t+1} \epsilon_1^* & & \uparrow 1 \otimes (\partial_{c-1}^0)^* \\
 \bigwedge^{t+1} F \otimes \bigwedge^t G^* \otimes S_0 G^* & \xleftarrow{\sigma_{c-2}^*} & G \otimes S_1 G^* \otimes \bigwedge^{t+1} F \otimes \bigwedge^t G^* \\
 \parallel & & \\
 (S_0 G \otimes \bigwedge^{c-2} F)^* & &
 \end{array}$$

We take $\{y_i\}_{i=1}^{t+c-1}$ to be a free R -basis of F and $\{y_i^*\}_{i=1}^{t+c-1}$ to be the dual basis, and we let $\{i_1, i_2, \dots, i_{t+1}\}$, $i_1 < i_2 < \dots < i_{t+1}$, be a subset of $I := \{1, 2, \dots, t+c-1\}$. According to [22] or [13, pp. 592–593] (see also the exact sequence (2.1)), ϵ_0^* and ϵ_1^* are defined by

$$\epsilon_0^*(g \otimes y_{i_1} \wedge y_{i_2} \wedge \dots \wedge y_{i_t}) := s_{t+1} \cdot g \otimes \varphi(y_{i_1}) \wedge \dots \wedge \varphi(y_{i_t})$$

and

$$\epsilon_1^*(y_{i_1} \wedge y_{i_2} \wedge \dots \wedge y_{i_{t+1}}) := \sum_{j=1}^{t+1} s_j \cdot (\varphi(y_{i_1}) \wedge \dots \wedge \varphi(y_{i_{j-1}}) \wedge \varphi(y_{i_{j+1}}) \wedge \dots \wedge \varphi(y_{i_{t+1}})) y_{i_j}$$

where s_j is the sign of the permutation of I that takes the elements of $\{i_1, i_2, \dots, i_{t+1}\} \setminus \{i_j\}$ into the first t positions. Note that $\varphi(y_i)$ are the columns of the matrix \mathcal{A} associated to φ , that $\varphi(y_{i_1}) \wedge \dots \wedge \varphi(y_{i_t})$ is the maximal minor corresponding to the columns i_1, \dots, i_t and that a replacement of ϵ_1^* by $(-1)^{t+1} \epsilon_1^*$, cf. the diagram above, still makes the leftmost column in Diagram A a free resolution of $S_{c-2}M$. We define

$$\sigma_c^* = \text{Id}_G$$

and

$$\sigma_{c-1}^*(g \otimes y_{i_1} \wedge \dots \wedge y_{i_t}) = \sum_{j=1}^t (\varphi(y_{i_1}) \wedge \dots \wedge \varphi(y_{i_{j-1}}) \wedge g \wedge \varphi(y_{i_{j+1}}) \wedge \dots \wedge \varphi(y_{i_t})) y_{i_j}.$$

A straightforward computation gives us the desired commutativity, namely,

$$\sigma_c^* \cdot \epsilon_0^* = \varphi \cdot \sigma_{c-1}^* \quad \text{and} \quad (-1)^{t+1} \epsilon_1^* \cdot \sigma_{c-2}^* = \sigma_{c-1}^* \cdot (1 \otimes (\partial_{c-1}^0)^*).$$

We will now define the morphism $\ell_\bullet : Q_\bullet \rightarrow F_\bullet[1]$. For any integer i , $0 \leq i \leq c-2$, we define

$$\ell_i : S_{c-2-i} G \otimes \bigwedge^i F \rightarrow F^* \otimes S_{c-2-i} G \otimes \bigwedge^{i+1} F$$

sending an element

$$m \otimes f \in S_{c-2-i}G \otimes \bigwedge^i F$$

to

$$\ell_i(m \otimes f) = \sum_{j=1}^{t+c-1} y_j^* \otimes m \otimes (y_j \wedge f) \in F^* \otimes S_{c-2-i}G \otimes \bigwedge^{i+1} F.$$

Using the diagram

$$\begin{array}{ccccc}
 & & \ell_{i+1} & & \\
 & \swarrow & & \searrow & \\
 S_{c-4-i}G \otimes \bigwedge^{i+2} F & \xrightarrow{\sigma_{i+2}} & G^* \otimes S_{c-3-i}G \otimes \bigwedge^{i+2} F & \xrightarrow{\varphi^* \otimes 1} & F^* \otimes S_{c-3-i}G \otimes \bigwedge^{i+2} F \\
 \downarrow \partial_{i+2}^{c-4-i} & & \downarrow 1 \otimes \partial_{i+2}^{c-3-i} & & \downarrow 1 \otimes \partial_{i+2}^{c-3-i} \\
 S_{c-3-i}G \otimes \bigwedge^{i+1} F & \xrightarrow{\sigma_{i+1}} & G^* \otimes S_{c-2-i}G \otimes \bigwedge^{i+1} F & \xrightarrow{\varphi^* \otimes 1} & F^* \otimes S_{c-2-i}G \otimes \bigwedge^{i+1} F \\
 \downarrow \partial_{i+1}^{c-3-i} & & \downarrow 1 \otimes \partial_{i+1}^{c-2-i} & & \downarrow 1 \otimes \partial_{i+1}^{c-2-i} \\
 S_{c-2-i}G \otimes \bigwedge^i F & \xrightarrow{\sigma_i} & G^* \otimes S_{c-1-i}G \otimes \bigwedge^i F & \xrightarrow{\varphi^* \otimes 1} & F^* \otimes S_{c-1-i}G \otimes \bigwedge^i F \\
 & & \ell_i & &
 \end{array}$$

and [13, Proposition A2.8, p. 583] onto the derivation $\varphi^*(x_j^*)$, we check for any i , $0 \leq i \leq c-3$, that

$$(\varphi^* \otimes 1) \cdot \sigma_{i+1} = (1_{F^*} \otimes \partial_{i+2}^{c-3-i}) \cdot \ell_{i+1} + \ell_i \cdot \partial_{i+1}^{c-3-i}.$$

We dualize the top part of Diagram A and define ℓ_c^* , ℓ_{c-1}^* (obviously $\ell_i^* = 0$ for $i \geq c+1$) as follows:

$$\ell_c^* = \text{Id}_F$$

and

$$\ell_{c-1}^*(f \otimes y_{i_1} \wedge \cdots \wedge y_{i_t}) = f \wedge y_{i_1} \wedge \cdots \wedge y_{i_t}.$$

A direct calculation using the following diagram:

$$\begin{array}{ccccc}
 & & \ell_c^* & & \\
 & \swarrow & & \searrow & \\
 0 & & 0 & & 0 \\
 \uparrow & & \uparrow & & \uparrow \\
 G & \xleftarrow{\sigma_c^*} & G \otimes R & \xleftarrow{\varphi \otimes 1} & F \otimes R \\
 \uparrow \varphi & & \uparrow 1 \otimes \epsilon_0^* & & \uparrow \epsilon_0^* \\
 F & \xleftarrow{\sigma_{c-1}^*} & G \otimes S_0 G^* \otimes \bigwedge^t F \otimes \bigwedge^t G^* & \xleftarrow{\varphi \otimes 1} & F \otimes S_0 G^* \otimes \bigwedge^t F \otimes \bigwedge^t G^* \\
 \uparrow (-1)^{t+1} \epsilon_1^* & & \uparrow 1 \otimes (\partial_{c-1}^0)^* & & \uparrow 1 \otimes (\partial_{c-1}^0)^* \\
 \bigwedge^{t+1} F \otimes \bigwedge^t G^* \otimes S_0 G & \xleftarrow{\sigma_{c-1}^*} & G \otimes S_1 G^* \otimes \bigwedge^{t+1} F \otimes \bigwedge^t G^* & \xleftarrow{\varphi \otimes 1} & F \otimes S_1 G^* \otimes \bigwedge^{t+1} F \otimes \bigwedge^t G^* \\
 & & \ell_{c-1}^* & &
 \end{array}$$

gives us (recall that $\sigma_c^* = 1$)

$$\varphi \cdot \ell_c^* = \sigma_c^* \cdot (\varphi \otimes 1)$$

and

$$(-1)^{t+1} \epsilon_1^* \cdot \ell_{c-1}^* + \ell_c^* \cdot \epsilon_0^* = \sigma_{c-1}^* \cdot (\varphi \otimes 1).$$

Now we are ready to apply Lemma 3.1. Since $\ell_c = \text{Id}$ and $\sigma_c = \text{Id}$, the corresponding summands split off and we deduce that $\text{pd Ext}_R^1(M, S_{c-1}M) = c$, i.e. $\text{Ext}_R^1(M, S_{c-1}M)$ is a maximal Cohen–Macaulay A -module.

Finally we prove the last assertion of the theorem. We assume that $a_j = 1$ for all j and $b_i = 0$ for all i . Recall that the degree of a linear standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension c is $\binom{t+c-1}{c}$. Using the exact sequence (3.3), we get

$$-2 \text{Ext}^1(M, S_{c-1}M) = 0$$

and

$$\begin{aligned} \dim_K(-1 \text{Ext}^1(M, S_{c-1}M)) &= (t+c-1) \cdot \binom{t+c-2}{c-1} \\ &= \text{rk}(\text{Ext}^1(M, S_{c-1}M)) \cdot \text{deg}(X). \end{aligned}$$

Therefore, $\text{Ext}^1(M, S_{c-1}M)$ is a maximal Cohen–Macaulay module maximally generated or, equivalently,

$$\widetilde{\text{Ext}_R^1(M, S_{c-1}M)}$$

is an Ulrich sheaf on X of rank c . □

Given an ACM scheme $X \subset \mathbb{P}^N$ with dualizing sheaf ω and a coherent sheaf \mathcal{E} on X , we denote by \mathcal{E}^ω the sheaf $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \omega)$. It is well known that \mathcal{E} is ACM if and only if \mathcal{E}^ω is ACM. So, as a Corollary of Theorem 3.2, we have:

Corollary 3.3. *We keep the notation introduced above, set $N := \text{Ext}_R^1(M, S_{c-1}M)$ and assume $\text{depth}_J A \geq 2$. Then:*

- $\text{Hom}_A(N, K_A)$ is a maximal Cohen–Macaulay A -module of rank c .
- If $a_j = 1$ for all j and $b_i = 0$ for all i , then $\widetilde{\text{Hom}_A(N, K_A)}$ is an Ulrich sheaf of rank c .

Remark 3.4. Keeping the notation introduced above and arguing as in the proof of Theorem 3.2, we can prove that for any i , $0 \leq i \leq c$, there exists an exact sequence

$$(3.6) \quad 0 \rightarrow S_{i-1}M \rightarrow G^* \otimes S_i M \rightarrow F^* \otimes S_i M \rightarrow \text{Ext}_R^1(M, S_i M) \rightarrow 0$$

provided $\text{depth}_J A \geq 2$ where $J = I_{t-1}(\mathcal{A})$. Note that we interpret $S_{-1}M$ as $\text{Hom}_A(M, A)$.

Hence we get a big diagram, similar to Diagram A above, where we have replaced $c-1$ by i . The part of the proof of Theorem 3.2 where we show the existence of $\{\sigma_j\}_{j=0}^c$ between the two leftmost columns (we call this part of the big diagram by (*)), seems to hold for any i , $0 \leq i \leq c$. Since σ_i is injective (except for $i = c$), this would imply that the length of the projective resolution of $\text{Ext}_R^1(M, S_i M)$ is at most $\text{pd } A + 1$ for $0 \leq i \leq c-1$. The only problem to get this result will be to define σ_j more generally and to verify the commutativity of the

first diagram in the proof of Theorem 3.2 where the dual of the “splice” maps ϵ_j , $j = 0, 1$, occur (we call this diagram (**)) after having replaced $c - 1$ by i and made the corresponding obvious changes).

The case $i = 1$ was treated in [24, proof of Theorem 3.1]. That proof is almost the dual of the proof of this part of Theorem 3.2. Indeed the lower part of diagram (*) for $i = 1$ is exactly diagram (**) provided we move G^* from the second column to G in the first column in diagram (*). Thus [24, proof of Theorem 3.1] implies this part of the proof of Theorem 3.2 and vice versa.

The existence of the morphisms σ_j in diagram (*) in the case $i = c$ is very similar (and easier) to what we had to prove in Theorem 3.2 for $i = c - 1$. Indeed, in this case we only need to check diagram (**) where now only one ϵ_j occur. The mapping cone construction leads to a resolution of $\text{Ext}_R^1(M, S_c M)$ where, however, the leftmost free module (of rank one) clearly does not split off, whence the length of a minimal resolution must be $\text{pd } A + 2$. As explained for $i = 1$ as almost the “dual” of $i = c - 1$ above, the case $i = 0$ is similarly “dual” to $i = c$. We deduce the existence of a morphism between the R -free resolutions of $S_{-1}M$ and $G^* \otimes A$. In this case it is easy to see that the leftmost free module in the resolution of $\text{Ext}_R^1(M, A)$ split off. In particular, we get

$$\text{pd Ext}_R^1(M, S_i M) \leq \text{pd } A + 1 \quad \text{for } i \in \{0, 1, c - 1\}.$$

The importance of proving $\text{pd Ext}_R^1(M, S_i M)$ to be small follows from our next proposition because it leads to good depth of “twisted normal modules”. Indeed, in [24] we use this to prove both conjectures appearing in [26] on the dimension and smoothness of the locus of determinantal schemes inside the Hilbert scheme.

Proposition 3.5. *With the above notation, set $J = I_{t-1}(\mathcal{A})$. We have:*

- (1) $\text{Ext}_R^1(M, S_{c-1}M) \cong \text{Hom}_A(I/I^2, S_{c-2}M)$ provided $\text{depth}_J A \geq 2$,
- (2) $\text{Ext}_R^1(M, S_i M) \cong \text{Hom}_A(I/I^2, S_{i-1}M)$, $0 \leq i \leq c$, $i \neq c - 1$, provided $\text{depth}_J A \geq 4$.

Proof. We prove (1) and (2) simultaneously. By Proposition 2.3(ii), the canonical module $K_A(v) \cong S_{c-1}(M)$ for some integer v and by Proposition 2.3(ii) M is a maximal Cohen–Macaulay A -module. Therefore, we have

$$(3.7) \quad \text{Ext}_A^i(M, K_A) = 0 \quad \text{for } i \geq 1.$$

Using (2.2) as in [24, proof of Theorem 4.1], we get that

$$(3.8) \quad \text{Ext}_A^j(M, S_i M) = 0 \quad \text{for } 1 \leq j \leq 2$$

provided $\text{depth}_J A \geq 4$.

Under the assumption $\text{depth}_J A \geq 2$, we have seen in (3.5) that for any i , $1 \leq i \leq c$,

$$\text{Hom}_A(M, S_i M) \cong S_{i-1}M$$

(true also for $i = 0$). In particular, we have

$$(3.9) \quad \text{Hom}_A(M, K_A) \cong S_{c-2}(M)(-v).$$

Notice that the isomorphism

$$\text{Hom}_R(M, S_i M) \cong \text{Hom}_A(M \otimes_R A, S_i M)$$

leads to a spectral sequence

$$\mathrm{Ext}_A^p(\mathrm{Tor}_q^R(M, A), S_i M) \Rightarrow \mathrm{Ext}_R^{p+q}(M, S_i M);$$

a part of the usual 5-term associated sequence is

$$\begin{aligned} 0 \rightarrow \mathrm{Ext}_A^1(M \otimes_R A, S_i M) &\rightarrow \mathrm{Ext}_R^1(M, S_i M) \rightarrow \mathrm{Hom}_A(\mathrm{Tor}_1^R(M, A), S_i M) \\ &\rightarrow \mathrm{Ext}_A^2(M \otimes_R A, S_i M), \end{aligned}$$

which using (3.7)–(3.8) and the exactness of the sequence

$$0 \longrightarrow \mathrm{Tor}_1^R(M, A) \longrightarrow M \otimes_R I \longrightarrow M \otimes_R R \xrightarrow{\cong} M \otimes_R A \longrightarrow 0$$

allows us to conclude that

$$\mathrm{Ext}_R^1(M, S_i M) \cong \mathrm{Hom}_A(M \otimes_R I, S_i M).$$

Using (2.2) we get

$$\begin{aligned} \mathrm{Hom}_A(M \otimes_R I, S_i M) &\cong \mathrm{Hom}_A(M \otimes I/I^2, S_i M) \\ &\cong \mathrm{Hom}_A(I/I^2, \mathrm{Hom}_A(M, S_i M)) \\ &\cong \mathrm{Hom}_A(I/I^2, S_{i-1} M), \end{aligned}$$

and we are done. \square

Theorem 3.6. *Let $X \subset \mathbb{P}^n$ be a standard determinantal scheme of codimension $c \geq 2$ associated to a $t \times (t + c - 1)$ matrix \mathcal{A} . Set $J = I_{t-1}(\mathcal{A})$ and assume $\mathrm{depth}_J R/I(X) \geq 2$. Then, $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M})$ is an ACM sheaf of rank c . In addition, if X is a linear standard determinantal scheme, then $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M})(-H)$ is an initialized Ulrich sheaf of rank c and it has a pure linear $\mathcal{O}_{\mathbb{P}^n}$ -resolution of the following type:*

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^n}(-c)^{a_c} \rightarrow \cdots \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1)^{a_1} \rightarrow \mathcal{O}_{\mathbb{P}^n}^{a_0} \rightarrow \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M})(-H) \rightarrow 0$$

with $a_0 = c \cdot \deg(X) = c \cdot \binom{t+c-1}{c}$ and $a_i = \binom{c}{i} \cdot a_0$ for $1 \leq i \leq c$. Finally, if X is a local complete intersection, then

$$\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M}) \cong \mathcal{N}_X \otimes \widetilde{S_{c-2}M}$$

and

$$H_*^i(X, \mathcal{N}_X \otimes \widetilde{S_{c-2}M}) = 0 \quad \text{for } 1 \leq i \leq n - c - 1.$$

Proof. We can apply Proposition 3.5 and we get

$$\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M}) \cong \widetilde{\mathrm{Ext}_R^1(M, S_{c-1}M)}.$$

Applying Theorem 3.2 we conclude that $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M})$ is an ACM sheaf of rank c on X and if X is a linear standard determinantal scheme, then $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M})(-H)$ is an initialized Ulrich sheaf of rank c , cf. Corollary 3.7 for the twist. In this case the minimal $\mathcal{O}_{\mathbb{P}^n}$ -resolution of $\mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_{c-2}M})(-H)$ is given by [15, Proposition 2.1]. \square

As an immediate application of Theorem 3.2 and its proof we obtain a free R -resolution of $\mathrm{Ext}_R^1(M, S_{c-1}M)$, and more generally of $\mathrm{Ext}_R^1(M, S_i M)$ for $0 \leq i \leq c$, which we make explicit for later use. In fact, we have the following corollary.

Corollary 3.7. *We keep the notation introduced above and call $Q_\bullet, P_\bullet, F_\bullet$ the minimal free R -resolutions of $S_{i-1}M, G^* \otimes S_i M, F^* \otimes S_i M$, respectively. Then $\text{Ext}_R^1(M, S_i M)$, for $0 \leq i \leq c$, has a free R -resolution of the following type:*

$$\begin{aligned} 0 \rightarrow Q_c \rightarrow Q_{c-1} \oplus P_c \rightarrow Q_{c-2} \oplus P_{c-1} \oplus F_c \rightarrow Q_{c-3} \oplus P_{c-2} \oplus F_{c-1} \\ \rightarrow \cdots \rightarrow Q_0 \oplus P_1 \oplus F_2 \rightarrow P_0 \oplus F_1 \rightarrow F_0 \rightarrow \text{Ext}_R^1(M, S_i M) \rightarrow 0 \end{aligned}$$

provided $\text{depth}_J A \geq 2$ where $J = I_{t-1}(\mathcal{A})$. In particular, we have

$$\text{depth Ext}_R^1(M, S_i M) \geq \text{depth } A - 2.$$

Moreover if $i = c - 1$ (resp. $i \in \{0, 1\}$), we may delete Q_c, P_c and Q_{c-1}, F_c (resp. Q_c and a subsummand of P_c) from this resolution, and we get

$$\text{depth Ext}_R^1(M, S_i M) \geq \text{depth } A - 1 \quad \text{for } i = 0, 1$$

and

$$\text{depth Ext}_R^1(M, S_{c-1}M) = \text{depth } A.$$

Proof. Using Lemma 3.1 and Remark 3.4, we get a free R -resolution of $\text{Ext}_R^1(M, S_i M)$ of the form above. If $i = c - 1$ (resp. $i \in \{0, 1\}$), the mentioned summands split off by the proof of Theorem 3.2 (resp. Remark 3.4). \square

Note that we do not claim that this resolution is minimal (not even for $i = c - 1$) since we have not carefully analyzed all possible cancelation of repeated direct summands in the mapping cone construction. Nevertheless, there is a particular case where we can assure that *all* repeated summands split off. Indeed, if $i = c - 1, a_j = 1$ for all j and $b_s = 0$ for all s , then the module $\text{Ext}_R^1(M, S_{c-1}M)$ is Ulrich. By Proposition 2.11 (ii), it has a pure linear resolution and therefore, we may delete not only Q_c, P_c and Q_{c-1}, F_c but also any other repeated summand split off.

We would like to know whether the rank c ACM (resp. Ulrich) sheaves on X constructed in the Theorem 3.2 are indecomposable. We will see in the next section that under some weak conditions \mathcal{N}_X and $\mathcal{H}om(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_i M})$, $0 \leq i \leq c - 1$, are in fact indecomposable (see Theorems 4.3 and 4.14).

Now we consider and discuss a Conjecture which generalizes the main result of this section. It is based on a series of examples computed with Macaulay2 ([19]):

Conjecture 3.8. Let $X \subset \mathbb{P}^n$ be a standard determinantal scheme of codimension $c \geq 2$ associated to a $t \times (t + c - 1)$ matrix \mathcal{A} . Set $I = I_t(\mathcal{A})$ and $J = I_{t-1}(\mathcal{A})$, and assume that $\text{depth}_J R/I \geq 2$. With the above notation, we conjecture that for all integers i with $0 \leq i \leq c$, $\text{Ext}_R^i(S_i M, S_{c-i} M)$ is an (indecomposable) maximal Cohen–Macaulay R/I -module of rank $\binom{c}{i}$. In addition, if X is a linear determinantal scheme, then $\text{Ext}_R^i(S_i M, S_{c-i} M)$ is an (indecomposable) Ulrich R/I -module of rank $\binom{c}{i}$.

Our next goal will be to use the spectral sequence

$$E_2^{p,q} = \text{Ext}_A^p(\text{Tor}_q^R(S_r M, A), S_s M) \Rightarrow \text{Ext}_R^{p+q}(S_r M, S_s M) \quad \text{for } -1 \leq s \leq c$$

to get some non-obvious isomorphic variations of the above conjecture. In fact, we have the following proposition.

Proposition 3.9. *With the above notation, if $\text{depth}_J A \geq i + 2$ and $-1 \leq s \leq s + e \leq c$, then*

$$\text{Ext}_R^p(S_r M, S_s M) \cong \text{Ext}_R^p(S_{r+e} M, S_{s+e} M) \quad \text{for any } p \leq i.$$

In particular, for $i \leq \frac{c+1}{2}$, we have

$$\text{Ext}_R^i(S_i M, S_{c-i} M) \cong \text{Ext}_R^i(A, S_{c-2i} M) \cong \text{Ext}_R^i(S_e M, S_{c-2i+e} M) \quad \text{for } 0 \leq e \leq 2i.$$

Proof. Since $T := \text{Spec}(A) \setminus V(J)$ is locally a complete intersection, the \mathcal{O}_T -modules

$$\widetilde{\text{Tor}}_q^R(\widetilde{S_r M}, \mathcal{O}_T) \cong \widetilde{\text{Tor}}_q^R(\mathcal{O}_T, \mathcal{O}_T) \otimes_{\mathcal{O}_T} \widetilde{S_r M}$$

and

$$\widetilde{\text{Tor}}_q^R(\mathcal{O}_T, \mathcal{O}_T) \cong \bigwedge^q (J/J^2)$$

are locally free on T , whence $\text{depth}_J S_s M \geq i + 2$ and (2.2) implies

$$\begin{aligned} (3.10) \quad E_2^{p,q} &= \text{Ext}_A^p(\text{Tor}_q^R(S_r M, A), S_s M) \\ &\cong \text{H}^p(T, \mathcal{H}om(\bigwedge^q (J/J^2) \otimes \widetilde{S_r M}, \widetilde{S_s M})) \\ &\cong \text{H}^p(T, \mathcal{H}om(\bigwedge^q (J/J^2), \widetilde{S_{s-r} M})) \quad \text{for } p \leq i. \end{aligned}$$

For the same reason, if $0 \leq e \in \mathbb{Z}$ and $s + e \leq c$, then $S_{s+e} M$ is MCM (see Proposition 2.3) and the diagram

$$\begin{array}{ccc} \text{Hom}_R(S_r M, S_s M) & \longrightarrow & \text{Hom}_R(S_{r+e} M, S_{s+e} M) \\ \downarrow \cong & & \downarrow \cong \\ \text{Hom}_A(S_r M \otimes_R A, S_s M) & \longrightarrow & \text{Hom}_A(S_{r+e} M \otimes_R A, S_{s+e} M) \end{array}$$

and (3.10) lead to isomorphisms

$$\text{Ext}_A^p(\text{Tor}_q^R(S_r M, A), S_s M) \cong \text{Ext}_A^p(\text{Tor}_q^R(S_{r+e} M, A), S_{s+e} M)$$

for any $p \leq i$, $-1 \leq s \leq s + e \leq c$. Hence, they converge to isomorphic Ext_R^p -groups and we get what we want. \square

Remark 3.10. Using Proposition 3.9, we can improve upon (2) of Proposition 3.5. Indeed, assuming $\text{depth}_J A \geq 3$, we get

$$\text{Ext}_R^1(M, S_i M) \cong \text{Ext}_R^1(S_{c-i} M, S_{c-1} M) \quad \text{for } -1 \leq i \leq c$$

by Proposition 3.9. We have $\text{Ext}_A^j(S_{c-i} M, S_{c-1} M) = 0$ for $j \geq 1$ because $S_{c-1} M$ is a twist of the canonical module and $S_{c-i} M$ is MCM. Since

$$\text{Hom}(\text{Tor}_1^R(S_{c-i} M, A), S_{c-1} M) \cong \text{Hom}_A(I/I^2, S_{i-1} M)$$

by (3.10), we get

$$\text{Ext}_R^1(M, S_i M) \cong \text{Hom}_A(I/I^2, S_{i-1} M)$$

under the assumption $\text{depth}_J A \geq 3$ by the spectral sequence of (3.10).

Now we will find some natural depth_J -conditions which will allow us to prove that

$$(3.11) \quad \text{Ext}_R^a(S_r M, S_s M) \cong \text{Hom}_A(\bigwedge^a (I/I^2), S_{s-r} M) \quad \text{for } 0 \leq a \leq r - s + c.$$

Lemma 3.11. *Let r, s and $a \geq 1$ be integers satisfying $-1 \leq s \leq c$ and assume*

$$(3.12) \quad \text{depth}_J \text{Hom}_A(\bigwedge^q (I/I^2), S_{s-r}M) \geq a + 3 - q \quad \text{for all } 0 \leq q < a.$$

Then

$$\text{Ext}_R^a(S_r M, S_s M) \cong \text{Hom}_A(\bigwedge^a (I/I^2), S_{s-r}M).$$

Remark 3.12. Inequality (3.12) for $q = 0$ just means $\text{depth}_J S_{s-r}M \geq a + 3$, whence we have $\text{depth}_J A \geq a + 3$. It follows that (3.10) holds for $p \leq a + 1$ provided $-1 \leq s \leq c$.

Proof. Using (3.10) and the hypothesis (3.12) which yields

$$(3.13) \quad \begin{aligned} & \text{H}^{a+1-q}(T, \mathcal{H}om(\bigwedge^q (\mathcal{J}/\mathcal{J}^2), \widetilde{S_{s-r}M})) \\ &= \text{H}_J^{a+2-q}(\text{Hom}(\bigwedge^q (I/I^2), S_{s-r}M)) = 0, \end{aligned}$$

we get

$$E_2^{p,q} = \text{Ext}_A^p(\text{Tor}_q^R(S_r M, A), S_s M) = 0 \quad \text{for } p + q = a + 1, 0 \leq q < a.$$

In the same way

$$\text{H}^{a-q}(T, \mathcal{H}om(\bigwedge^q (\mathcal{J}/\mathcal{J}^2), \widetilde{S_{s-r}M})) = 0$$

implies that

$$E_2^{p,q} = 0 \quad \text{for } p + q = a, 0 \leq q < a.$$

Hence all terms $E_2^{p,q}$ with $p + q = a$ vanish except $E_2^{0,a}$ and we get

$$\text{Ext}_R^a(S_r M, S_s M) \cong E_\infty^{0,a}.$$

We claim that (3.13) implies $E_\infty^{0,a} \cong E_2^{0,a}$. Indeed, by (3.13), we have $E_\mu^{p,q} = 0$ for any $\mu \geq 2$ and (p, q) satisfying $p + q = a + 1, 0 \leq q < a$. Therefore, the differentials of the spectral sequence

$$d_{\mu, 1-\mu} : E_\mu^{0,a} \rightarrow E_\mu^{\mu, a+1-\mu}, \quad \mu \geq 2,$$

vanish for $\mu \geq 2$ because $E_\mu^{\mu, a+1-\mu} = 0$ for $\mu \geq 2$. It follows that

$$(3.14) \quad \text{Ext}_R^a(S_r M, S_s M) \cong E_2^{0,a} \cong \text{Hom}_A(\bigwedge^a (I/I^2), S_{s-r}M)$$

where the isomorphism to the right follows from $\text{depth}_J S_{s-r}M \geq 2$. \square

Example 3.13. Suppose $\text{depth}_J A \geq 5$ and take $a = r = 2$. Then (3.12) is satisfied for $s = 2$; the case $q = 0$ by hypothesis and the case $q = 1$ follows from [24, Theorem 5.11]. Therefore, we get

$$(3.15) \quad \text{Ext}_R^2(S_2 M, S_2 M) \cong \text{Hom}_A(\bigwedge^2 I/I^2, A).$$

If $s = c$, then (3.12) is satisfied; the case $q = 1$ follows from Theorem 3.2 and Proposition 3.5, whence

$$\text{Ext}_R^2(S_2 M, S_c M) \cong \text{Hom}_A(\bigwedge^2 I/I^2, S_{c-2}M).$$

Lemma 3.14. *Let $1 \leq r \leq c$ and suppose $\text{depth}_J A \geq r + 1$. Then for any integer s with $r - 1 \leq s \leq c$ we have*

$$(3.16) \quad \text{pd Ext}_R^r(S_r M, S_s M) \leq \text{pd } A + 2r.$$

Proof. We proceed by induction on r . The case $r = 1$ follows from Corollary 3.7. Suppose

$$\text{pd Ext}_R^i(S_i M, S_{s-r+i} M) \leq \text{pd } A + 2i \quad \text{for } 1 \leq i < r.$$

First note that by Proposition 3.9 we have

$$\text{Ext}_R^i(S_i M, S_{s-r+i} M) \cong \text{Ext}_R^i(S_r M, S_s M).$$

By Proposition 2.3, we have a resolution of length c of $S_r M$ of the following type:

$$\begin{aligned} \rightarrow L_{r+1} \xrightarrow{\epsilon} L_r = \bigwedge^r F \rightarrow L_{r-1} = G \otimes \bigwedge^{r-1} F \\ \rightarrow \dots \rightarrow L_1 = S_{r-1} G \otimes F \rightarrow L_0 = S_r G \rightarrow S_r M \rightarrow 0 \end{aligned}$$

where ϵ is the ‘‘splice map’’. Applying the contravariant functor $\text{Hom}_R(-, S_s M)$, we get a complex

$$(3.17) \quad \begin{aligned} 0 \longrightarrow S_s M \otimes L_0^* \xrightarrow{\delta_0} S_s M \otimes L_1^* \xrightarrow{\delta_1} \dots \xrightarrow{\delta_{r-2}} S_s M \otimes L_{r-1}^* \\ \xrightarrow{\delta_{r-1}} S_s M \otimes L_r^* \xrightarrow{\epsilon^*} S_s M \otimes L_{r+1}^* \end{aligned}$$

where $\ker(\delta_0) = \text{Hom}(S_r M, S_s M) \cong S_{s-r} M$, $\epsilon^* = \text{Hom}(\epsilon, S_s M) = 0$, whence

$$\text{Ext}_R^r(S_r M, S_s M) = S_s M \otimes L_r^* / \text{im}(\delta_{r-1}).$$

Due to the horseshoe lemma and the exact sequence

$$0 \rightarrow \text{im}(\delta_{i-1}) \rightarrow \ker(\delta_i) \rightarrow \text{Ext}_R^i(S_r M, S_s M) \rightarrow 0,$$

we get the implication

$$\text{pd im}(\delta_{i-1}) \leq \text{pd } A + 2i - 1 \implies \text{pd ker}(\delta_i) \leq \text{pd } A + 2i.$$

The exact sequence

$$0 \rightarrow \ker(\delta_i) \rightarrow S_s M \otimes L_i^* \rightarrow \text{im}(\delta_i) \rightarrow 0$$

shows the implication

$$\text{pd ker}(\delta_i) \leq \text{pd } A + 2i \implies \text{pd im}(\delta_i) \leq \text{pd } A + 2i + 1 \quad \text{for } 1 \leq i < r.$$

Since by (3.17) and $\ker(\delta_0) \cong S_{s-r} M$ we have

$$\text{pd im}(\delta_0) \leq \text{pd } A + 1,$$

we conclude that $\text{pd im}(\delta_{r-1}) \leq \text{pd } A + 2r - 1$, and using the exact sequence

$$0 \rightarrow \text{im}(\delta_{r-1}) \rightarrow S_r(M) \otimes L_r^* \rightarrow \text{Ext}_R^r(S_r M, S_s M) \rightarrow 0,$$

we get

$$\text{pd Ext}_R^r(S_r M, S_s M) \leq \text{pd } A + 2r$$

which proves the lemma. \square

Remark 3.15. If $s \in \{r - 1, r, r + c - 2\}$, $s \leq c$, we have

$$\text{pd Ext}_R^r(S_r M, S_s M) \leq \text{pd } A + 1 \quad \text{for } r = 1$$

by Remark 3.4. Using the proof above, we show $\text{pd ker}(\delta_1) \leq \text{pd } A + 1$, whence we can improve upon the conclusion and we get

$$\text{pd Ext}_R^r(S_r M, S_s M) \leq \text{pd } A + 2r - 1 \quad \text{for } 1 \leq r \leq c.$$

Theorem 3.16. *Let $X \subset \mathbb{P}^n$ be a standard determinantal scheme of codimension $c \geq 2$ associated to a $t \times (t + c - 1)$ matrix \mathcal{A} . Let $I = I_t(\mathcal{A})$, $J = I_{t-1}(\mathcal{A})$ and $A = R/I$. Fix two integers r, s with $r - 1 \leq s \leq c$ and assume that $\text{depth}_J A \geq 2a + 2$ (resp. $\text{depth}_J A \geq 2a + 1$ if $s \in \{r - 1, r, r + c - 2\}$ and $a > 1$). Then, we have*

$$\text{Ext}_R^a(S_r M, S_s M) \cong \text{Hom}_A\left(\bigwedge^a(I/I^2), S_{s-r} M\right) \quad \text{for } 0 \leq a \leq r - s + c.$$

Remark 3.17. It is worthwhile to point out that Theorem 3.16 with $r = s = 0$ holds more generally in the licci case (see [28, Proposition 13 and Remark 14]).

Proof. Indeed, since $-1 \leq s - r + a \leq c$, we can apply Proposition 3.9 and we have

$$\text{Ext}_R^a(S_r M, S_s M) \cong \text{Ext}_R^a(S_a M, S_{s-r+a} M).$$

Now, we apply Lemma 3.14 and Auslander–Buchsbaum’s formula to deduce that

$$(3.18) \quad \text{depth}_J \text{Ext}_R^a(S_r M, S_s M) \geq \text{depth}_J A - 2a$$

(true also for $a = 0$ since we may assume $r - s + c \geq 0$ in which case $S_{s-r} M$ is MCM).

Let q be an integer such that $0 \leq q < a$. Then (3.18) holds replacing a by q . Since we may by induction suppose that

$$(3.19) \quad \text{Ext}_R^q(S_r M, S_s M) \cong \text{Hom}_A\left(\bigwedge^q(I/I^2), S_{s-r} M\right),$$

it follows by assumption that

$$(3.20) \quad \text{depth}_J \text{Hom}_A\left(\bigwedge^q(I/I^2), S_{s-r} M\right) \geq \text{depth}_J A - 2q \geq 2a + 2 - 2q \geq a + 3 - q;$$

and applying Lemma 3.11 we get the result for $s \notin \{r - 1, r, r + c - 2\}$.

Finally if $s \in \{r - 1, r, r + c - 2\}$ we use Remark 3.15 to improve upon (3.18) for $a > 0$ and hence upon (3.19) for $q > 0$. Using $\text{depth}_J A \geq 2a + 1$, we still get (3.20), and we are done. \square

Corollary 3.18. *Let $i \geq 1$, $s \geq -1$ and a be integers satisfying $0 \leq a \leq c - s$.*

- (i) *Suppose that $\text{depth}_J A \geq 2a + 2 + i$ (resp. $\text{depth}_J A \geq 2a + 1 + i$ if $s \in \{-1, 0, c - 2\}$ and $a > 0$). Then, for $1 \leq j \leq i$, we have*

$$\text{Ext}_A^j\left(\bigwedge^a(I/I^2), S_s M\right) = 0 \quad \text{and} \quad H_*^j(X \setminus V(J), \left(\bigwedge^a \mathcal{I}_X / \mathcal{I}_X^2\right)^\vee \otimes \widehat{S_s M}) = 0.$$

- (ii) *Suppose that $\dim A \geq 2a + 2 + i$ (resp. $\geq 2a + 1 + i$ if $s \in \{-1, 0, c - 2\}$ and $a > 0$). Then we have*

$$H_*^j(X, \mathcal{H}om\left(\bigwedge^a(\mathcal{I}_X / \mathcal{I}_X^2), \widehat{S_s M}\right)) = 0 \quad \text{for } 1 \leq j \leq i.$$

Proof. (i) Taking $r = 0$ in Theorem 3.16, we get

$$\text{Ext}_R^a(A, S_s M) \cong \text{Hom}_A\left(\bigwedge^a(I/I^2), S_s M\right)$$

and applying Lemma 3.14 together with Auslander–Buchsbaum’s formula, as in (3.18), we have

$$(3.21) \quad \text{depth}_J \text{Hom}_A\left(\bigwedge^a(I/I^2), S_s M\right) \geq 2 + i.$$

Since $\bigwedge^a(\mathcal{J}_X/\mathcal{J}_X^2)$ is locally free on $X \setminus V(J)$, we apply (2.2) and, for $1 \leq j \leq i$, we get

$$\text{Ext}_A^j(\bigwedge^a(I/I^2), S_s M) = 0$$

and

$$H_*^j(X \setminus V(J), \mathcal{H}om(\bigwedge^a(\mathcal{J}_X/\mathcal{J}_X^2), \widetilde{S_s M})) = 0.$$

Suppose $s \in \{-1, 0, c - 2\}$. Then, for $a = 1$, we may use Remark 3.15 to improve upon (3.18) by 1 and since we still get (3.21), we conclude by the arguments above. If $a \geq 2$, then Theorem 3.16 applies, and using Remark 3.15 to improve upon (3.18), we get (3.21) and we conclude as previously.

(ii) The argument using the projective dimension and Auslander–Buchsbaum’s formula also imply (3.21) with \mathfrak{m} instead of J , whence we get $H_{\mathfrak{m}}^{j+1}(\text{Hom}_A(\bigwedge^a(I/I^2), S_s M)) = 0$ and we are done. \square

The case $a = 1, i = 1$ and $s = -1$ of Corollary 3.18 is of special interest because it shows the vanishing of a group that we tried very much to compute in [25]. Indeed, we have

Corollary 3.19. *Let $J = I_{t-1}(\mathcal{A})$ and $A = R/I$ and suppose $\text{depth}_J A \geq 4$. Then*

$$\text{Ext}_A^1(I/I^2, \text{Hom}(M, A)) = 0.$$

In [25, Theorem 5.1] we repeatedly need to apply Corollary 3.19 to standard determinantal schemes obtained by deleting at least one column. Indeed, Corollary 3.19 shows that all assumptions of [25, Theorem 5.1 (ii)] are satisfied in the case $\dim X \geq 2, a_j \geq b_i$ for any i, j . It follows that the closure of the locus of determinantal schemes W inside the Hilbert scheme is a generically smooth component of the Hilbert scheme (if $W \neq \emptyset$)! Thus Corollary 3.19 partially reproves [24, Corollary 5.9]. In particular, we also get [26, Conjecture 4.2] from [25, Theorem 5.1] and Corollary 3.19.

Since the case $a = 1$ and $s = c - 1$ is related to the dual of the conormal module of A , we remark

Corollary 3.20. *Let $J = I_{t-1}(\mathcal{A}), A = R/I, X = \text{Proj}(A)$ and let K_A be the canonical module of A . If $\text{depth}_J A \geq 4 + i$, then*

$$H_{\mathfrak{m}}^k(I/I^2) = 0 \quad \text{for } \dim X - i < k \leq \dim X,$$

or equivalently

$$\text{Ext}_A^j(I/I^2, K_A) = 0 \quad \text{for } 1 \leq j \leq i.$$

Proof. This is immediate from Corollary 3.18 (i) since $K_A(v) \cong S_{c-1}(M)$ for some integer v by Proposition 2.3 (ii). For the equivalent statement, we use Gorenstein duality. \square

As another application of Theorem 3.16, we can partially restate Conjecture 3.8. Indeed,

Conjecture 3.21. Let $X \subset \mathbb{P}^n$ be a linear standard determinantal scheme of codimension $c \geq 2$ associated to a $t \times (t + c - 1)$ matrix \mathcal{A} . Set $I = I_t(\mathcal{A})$ and $J = I_{t-1}(\mathcal{A})$. For any integer $a, 0 \leq a \leq \frac{c+1}{2}$, we conjecture that $\text{Hom}(\bigwedge^a(I/I^2), S_{c-2a}M)$ is an (indecomposable) Ulrich R/I -module of rank $\binom{c}{a}$ provided $\text{depth}_J A \geq 2a + 2$.

Notice that under the above hypothesis Conjectures 3.8 and 3.21 are equivalent since, by Theorem 3.16, we have

$$\mathrm{Ext}_R^a(S_a M, S_{c-a} M) \cong \mathrm{Hom}_A\left(\bigwedge^a (I/I^2), S_{c-2a} M\right) \quad \text{for } 1 \leq a \leq \frac{c+1}{2}.$$

We want to point out that for $a = 0$, the conjecture was proved in [4, Proposition 2.8] and in this section, we prove it for $a = 1$ (and, indecomposability for $n \geq 2c + 1$ in the next section).

4. The indecomposability of the normal sheaf of a determinantal variety

In this section we address Problem 1.1 (3) and we determine conditions under which the normal sheaf \mathcal{N}_X , and more generally the “twisted” normal sheaves

$$\mathcal{N}_X(\mathcal{M}^i) := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S_i M}), \quad -1 \leq i \leq c-1,$$

of a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension c are simple, i.e.

$$\mathrm{Hom}(\mathcal{N}_X(\mathcal{M}^i), \mathcal{N}_X(\mathcal{M}^i)) \cong K,$$

and, hence, indecomposable (cf. Theorems 4.3 and 4.14). Obviously $\mathcal{N}_X(\mathcal{M}^0) = \mathcal{N}_X$. We keep the notation of previous sections. So, \mathcal{A} will be the $t \times (t + c - 1)$ matrix associated to the standard determinantal scheme $X \subset \mathbb{P}^n$, $I = I_t(\mathcal{A})$, $A = R/I$ and

$$(4.1) \quad \cdots \rightarrow \bigoplus_j R(-n_{2j}) \rightarrow \bigoplus_i R(-n_{1i}) \rightarrow R \rightarrow A \rightarrow 0$$

the Eagon–Northcott resolution of A . In this section we often have $d_{i,j} := a_j - bi > 0$ for all i, j .

Again we assume $c > 1$, since the case $c = 1$ corresponds to a hypersurface $X \subset \mathbb{P}^n$ and $\mathcal{N}_X \cong \mathcal{O}_X(\delta)$, $\delta := \deg(X)$, is simple. We also assume $t > 1$, since the case $t = 1$ corresponds to a codimension c complete intersection $X \subset \mathbb{P}^n$ and $\mathcal{N}_X(\mathcal{M}^i)$, $\mathcal{N}_X = \bigoplus_{i=1}^c \mathcal{O}_X(d_i)$, $d_i \in \mathbb{Z}$, are neither simple nor indecomposable. Let us start computing the depth of the conormal bundle of a standard determinantal scheme. We have:

Proposition 4.1. *Let \mathcal{A} be a $t \times (t + c - 1)$ homogeneous matrix with entries that are general forms of degree $d_{ij} > 0$, let $I = I_t(\mathcal{A})$, $J = I_{t-1}(\mathcal{A})$ and $X \subset \mathbb{P}^n$ be the standard determinantal scheme of codimension c associated to \mathcal{A} . Assume that $n \geq 2c - 2$. Then, it holds:*

- (1) $\mathrm{depth}_{\mathfrak{m}} I/I^2 = n - 2c + 2$,
- (2) $\mathrm{depth}_J I/I^2 \geq 3$ (resp. $= n - 2c + 2$) provided $n \geq 2c + 1$ (resp. $n \leq 2c$).

Therefore, if $n \geq 2c$, there is a closed subset $Z \subset X$ such that $X \setminus Z \hookrightarrow \mathbb{P}^n$ is a local complete intersection and $\mathrm{depth}_{I(Z)} I/I^2 \geq 2$.

Proof. (1) We will first prove Proposition 4.1 for an ideal generated by the maximal minors of a matrix with entries that are indeterminates and we will deduce that it also works for a homogeneous matrix with entries of general homogeneous polynomials.

So, we first assume that $n = t(t + c - 1) - 1$ and \mathcal{A} is a matrix with entries x_0, x_1, \dots, x_n of indeterminates. Set $R = k[x_0, \dots, x_n]$, $I = I_t(\mathcal{A})$ and call Ω the module of differentials of R/I over k . By [6, Theorem 14.12],

$$\text{depth}_{\mathfrak{m}} \Omega = \text{depth}_{\mathfrak{m}} R/I - \text{depth}_J R/I + 2$$

and note that $\text{depth}_J R/I = \text{codim}(I_{t-1}(\mathcal{A}), R/I)$. Since

$$\text{depth}_{\mathfrak{m}} R/I = \dim R/I = n - c + 1$$

and

$$\text{codim}(I_{t-1}(\mathcal{A}), R/I) = \text{codim}(I_{t-1}(\mathcal{A})) - \text{codim}(I_t(\mathcal{A})) = 2(c + 1) - c = c + 2,$$

we get

$$\text{depth}_{\mathfrak{m}} \Omega = (n - c + 1) - (c + 2) + 2 = t(t + c - 1) - 2c.$$

Therefore, using the exact sequence

$$0 \rightarrow I/I^{(2)} \rightarrow (R/I)^{t(t+c-1)} \rightarrow \Omega \rightarrow 0,$$

we deduce that

$$\begin{aligned} \text{depth}_{\mathfrak{m}} I/I^{(2)} &= \text{depth}_{\mathfrak{m}} \Omega + 1 \\ &= t(t + c - 1) - 2c + 1 \\ &= n - 2c + 2. \end{aligned}$$

By [6, Corollary 9.18], we have $I^2 = I^{(2)}$. Therefore, we conclude that

$$\text{depth}_{\mathfrak{m}} I/I^2 = n - 2c + 2.$$

Let us now assume $n < t(t + c - 1) - 1$. We distinguish two cases:

Case 1: Assume $d_{ij} = 1$ for all i, j (i.e. the entries of the matrix \mathcal{A} are linear forms). We have a $t \times (t + c - 1)$ matrix $\overline{\mathcal{A}} = (x_{i,j})$ of indeterminates and the ideal $I_t(\overline{\mathcal{A}}) \subset S := k[x_{i,j}]$ which verifies

$$\text{depth}_{\mathfrak{m}_S} I_t(\overline{\mathcal{A}})/I_t(\overline{\mathcal{A}})^2 = t(t + c - 1) - 2c + 1.$$

We choose $t(t + c - 1) - n - 1$ general linear forms $\ell_1, \dots, \ell_{t(t+c-1)-n-1} \in S = k[x_{i,j}]$ and we set

$$S/(\ell_1, \dots, \ell_{t(t+c-1)-n-1}) \cong k[x_0, x_1, \dots, x_n] =: R.$$

Let us call $I \subset R$ the ideal of R isomorphic to the ideal $I_t(\overline{\mathcal{A}})/(\ell_1, \dots, \ell_{t(t+c-1)-n-1})$ of $S/(\ell_1, \dots, \ell_{t(t+c-1)-n-1})$. Notice that I is nothing but the ideal $I_t(\mathcal{A})$ where $\mathcal{A} = (m_{i,j})$ is a $t \times (t + c - 1)$ homogeneous matrix with entries that are linear forms in $k[x_0, x_1, \dots, x_n]$ obtained from $\overline{\mathcal{A}} = (x_{i,j})$ by substituting using the equations $\ell_1, \dots, \ell_{t(t+c-1)-n-1}$. Since

$$\text{depth}_{\mathfrak{m}_S} I_t(\overline{\mathcal{A}})/I_t(\overline{\mathcal{A}})^2 = t(t + c - 1) - 2c + 1$$

and

$$t(t + c - 1) - n - 1 \leq t(t + c - 1) - 2c + 1,$$

we can assume that $\ell_1, \dots, \ell_{t(t+c-1)-n-1}$ is a regular sequence on both $I_t(\overline{\mathcal{A}})/I_t(\overline{\mathcal{A}})^2$ and $S/I_t(\overline{\mathcal{A}})$; and we conclude that

$$\text{depth}_{\mathfrak{m}_R} I/I^2 = \text{depth}_{\mathfrak{m}_S} I_t(\overline{\mathcal{A}})/I_t(\overline{\mathcal{A}})^2 - (t(t + c - 1) - n - 1) = n - 2c + 2.$$

Case 2: Since $d_{i,j} > 0$ for all i, j , it is enough to raise the entry $m_{i,j}$ of the above matrix \mathcal{A} to the power $d_{i,j}$.

(2) It follows from (1) and

$$\text{depth}_J I/I^2 \geq \text{depth}_J A - (\dim A - \text{depth}_{\mathfrak{m}} I/I^2),$$

cf. [23, Lemma 7] because $\text{depth}_J A = c+2$ (resp. $n+1-c$) for $n \geq 2c+1$ (resp. $n \leq 2c$). \square

As an immediate and nice consequence of the above result, we get for $n \geq 2c+1$ that the cohomology of the conormal bundle $H_{\mathfrak{m}}^j(I/I^2)$ is non-zero for only one value of $j \neq \dim R/I$, cf. [1] for $c=2$. Analogous result for the normal bundle was proved by Kleppe in [24, Theorem 5.11].

Corollary 4.2. *Let \mathcal{A} be a $t \times (t+c-1)$ homogeneous matrix with entries that are general forms of positive degree. Set $J = I_{t-1}(\mathcal{A})$, $A = R/I$ and let K_A be the canonical module of A .*

- (i) *We have $H_{\mathfrak{m}}^k(I/I^2) = 0$ for $k < n - 2c + 2$ or, equivalently, $\text{Ext}_A^j(I/I^2, K_A) = 0$ for $c \leq j \leq n - c + 1$.*
- (ii) *We have $H_{\mathfrak{m}}^k(I/I^2) = 0$ for $\max(3, n - 2c + 2) < k < n - c + 1$.*

Proof. (i) It follows from Proposition 4.1 and Gorenstein duality.

(ii) Since $\text{depth}_J A = \min(c+2, n-c+1)$, we can apply Corollary 3.20 and we get that $H_{\mathfrak{m}}^k(I/I^2) = 0$ for $n - 2c + 2 < k \leq \dim X = n - c$. Combining with (i) we get what we want. \square

Theorem 4.3. *Let $X \subset \mathbb{P}^n$ be a standard determinantal scheme of codimension $c \geq 2$ associated to a $t \times (t+c-1)$ matrix \mathcal{A} with entries that are general forms of positive degree. Let $\mathcal{N}_X(\mathcal{M}^i) := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S}_i \mathcal{M})$ for $-1 \leq i \leq c-1$ and assume $n \geq 2c$. Then,*

$$\text{Hom}(\mathcal{N}_X(\mathcal{M}^i), \mathcal{N}_X(\mathcal{M}^i)) \cong {}_0\text{Hom}_A(I/I^2, I/I^2) \cong K$$

provided $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$. In particular, \mathcal{N}_X and $\mathcal{N}_X(\mathcal{M}^i)$ are simple, and thus indecomposable.

Proof. First of all we observe that $\text{Hom}_R(I, I) \cong R$. Indeed, we have the following diagram:

$$\begin{array}{ccccccc} & & & 0 & & & \\ & & & \downarrow & & & \\ & & & \text{Hom}_R(R, R) = R & & & \\ & & & \downarrow & & & \\ 0 & \longrightarrow & \text{Hom}_R(I, I) & \longrightarrow & \text{Hom}_R(I, R) & \longrightarrow & \text{Hom}_R(I, A) \longrightarrow \text{Ext}_R^1(I, I) \\ & & & & \downarrow & & \\ & & & & \text{Ext}_R^1(A, R) = 0 & & \end{array}$$

Since $c \geq 2$, we have

$$\mathrm{Ext}_R^1(A, R) = 0 \quad \text{and} \quad R \cong \mathrm{Hom}_R(R, R) \cong \mathrm{Hom}_R(I, R).$$

Hence, it follows that $\mathrm{Hom}_R(I, I)$ is an ideal of $\mathrm{Hom}_R(I, R) = R$ containing the identity; and so $\mathrm{Hom}_R(I, I) \cong R$.

Claim. *If $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$, then $K \cong {}_0\mathrm{Hom}_R(I, I) \cong {}_0\mathrm{Hom}_A(I/I^2, I/I^2)$.*

Proof of the Claim. We apply $\mathrm{Hom}_R(-, I)$ and $\mathrm{Hom}_R(-, I/I^2)$ to the minimal resolution of I deduced from (4.1) and we get the following commutative diagram with exact horizontal sequences

$$\begin{array}{ccccccc}
 & & & 0 & & 0 & \\
 & & & \uparrow & & \uparrow & \\
 0 & \longrightarrow & \mathrm{Hom}_R(I, I/I^2) & \longrightarrow & \bigoplus_i I/I^2(n_{1i}) & \longrightarrow & \bigoplus_j I/I^2(n_{2j}) \longrightarrow \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & \mathrm{Hom}_R(I, I) & \longrightarrow & \bigoplus_i I(n_{1i}) & \longrightarrow & \bigoplus_j I(n_{2j}) \longrightarrow \\
 & & & & \uparrow & & \uparrow \\
 & & & & \bigoplus_i I^2(n_{1i}) & \longrightarrow & \bigoplus_j I^2(n_{2j}) \longrightarrow \\
 & & & & \uparrow & & \uparrow \\
 & & & & 0 & & 0.
 \end{array}$$

Since $\mathrm{Hom}_R(I, I/I^2) \cong \mathrm{Hom}_A(I/I^2, I/I^2)$ and ${}_0\mathrm{Hom}_R(I, I^2) = 0$, it is sufficient to show that $(I^2(n_{2j}))_0 = 0$ for all j . Using the natural surjective map $S_2I \twoheadrightarrow I^2$, it suffices to show that $(S_2I)_\mu = 0$ for $\mu := \max\{n_{2j}\}$. But $(S_2I)_\mu = 0$ because we have a surjective map

$$\bigoplus_{i \leq j} R(-n_{1i} - n_{1j})_\mu \cong S_2 \left(\bigoplus_i R(-n_{1i}) \right)_\mu \twoheadrightarrow (S_2I)_\mu$$

and

$$\bigoplus_{i \leq j} R(-n_{1i} - n_{1j})_\mu = 0$$

by the assumption $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$. Hence, the claim is proved. \square

Let us now prove that $\mathcal{N}_X(\mathcal{M}^i)$ is simple, i.e.

$$\mathrm{Hom}(\mathcal{N}_X(\mathcal{M}^i), \mathcal{N}_X(\mathcal{M}^i)) \cong {}_0\mathrm{Hom}_A(I/I^2, I/I^2) \cong K.$$

Set $J = I_{t-1}(\mathcal{A})$. Since $\mathrm{depth}_J R/I = \min(c+2, n+1-c) \geq 2$, we get that

$$\mathrm{depth}_J N_M \geq 2$$

where $N_M := \mathrm{Hom}_A(I/I^2, S_i M)$. This also implies $\mathrm{depth}_J \mathrm{Hom}_A(N_M, N_M) \geq 2$, whence

$$\begin{aligned}
 \mathrm{Hom}_A(N_M, N_M) &\cong \mathrm{H}_*^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{N}_X(\mathcal{M}^i), \mathcal{N}_X(\mathcal{M}^i))) \\
 &\cong \mathrm{H}_*^0(X, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{N}_X(\mathcal{M}^i), \mathcal{N}_X(\mathcal{M}^i)))
 \end{aligned}$$

where $Z := V(J)$. Using that $\mathcal{N}_X(\mathcal{M}^i) \cong (\mathcal{J}_X/\mathcal{J}_X^2)^\vee \otimes \widetilde{S}_i \widetilde{M}$ is locally free on $X \setminus Z$, we have

$$H_*^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{N}_X(\mathcal{M}^i), \mathcal{N}_X(\mathcal{M}^i))) = H_*^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{J}_X/\mathcal{J}_X^2, \mathcal{J}_X/\mathcal{J}_X^2)).$$

Since $n \geq 2c$, we can apply Proposition 4.1 and we get $\text{depth}_{I(Z)} I/I^2 \geq 2$ which implies that $\text{depth}_{I(Z)} \text{Hom}_A(I/I^2, I/I^2) \geq 2$, whence

$$\text{Hom}_A(I/I^2, I/I^2) \cong H_*^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{J}_X/\mathcal{J}_X^2, \mathcal{J}_X/\mathcal{J}_X^2)).$$

Putting altogether we obtain

$$H_*^0(X, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{N}_X(\mathcal{M}^i), \mathcal{N}_X(\mathcal{M}^i))) \cong \text{Hom}_A(I/I^2, I/I^2);$$

and taking the degree zero piece of these graded modules we get what we want. \square

Remark 4.4. In Theorem 4.3, the hypothesis $n \geq 2c$ together with the generality of the entries can be replaced by

$$(4.2) \quad H_{I(Z)}^1(I/I^2)_{n_{1i}} = H_{I(Z)}^0(I/I^2)_{n_{1i}} = H_{I(Z)}^0(I/I^2)_{n_{2j}} = 0 \quad \text{for any } i \text{ and } j$$

where $Z \subset X$ is a closed subset such that $X \setminus Z \hookrightarrow \mathbb{P}^n$ is a local complete intersection. Note that $I(Z) = \mathfrak{m}$ if we can take $Z = \emptyset$, e.g. if $n \leq 2c + 1$ and the entries are general forms. To show it, observe that the exact cohomology sequence associated to $0 \rightarrow I^2 \rightarrow I \rightarrow I/I^2 \rightarrow 0$ gives us ($Z := V(J)$)

$$H_{I(Z)}^1(I/I^2)_\mu \cong H_{I(Z)}^2(I^2)_\mu \cong H^1(X \setminus Z, \mathcal{J}_X^2(\mu))$$

and

$$H_{I(Z)}^0(I/I^2)_{n_{2i}} \cong H_{I(Z)}^1(I^2)_{n_{2i}} \cong H^0(X \setminus Z, \mathcal{J}_X^2(n_{2i}))$$

since $I^2(n_{2i})_0 = 0$. In the proof of Theorem 4.3, we show

$$\text{Hom}_A(N_M, N_M) \cong H_*^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{J}_X/\mathcal{J}_X^2, \mathcal{J}_X/\mathcal{J}_X^2))$$

and we use the hypothesis $n \geq 2c$ to get $\text{depth}_{I(Z)} I/I^2 \geq 2$ and, hence,

$$(4.3) \quad H_*^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{J}_X/\mathcal{J}_X^2, \mathcal{J}_X/\mathcal{J}_X^2)) \cong \text{Hom}_A(I/I^2, I/I^2).$$

To get Theorem 4.3 we only need the isomorphism (4.3) in degree 0. Letting $\text{Ext}_{I(Z)}^j(N, -)$ be the right derived functor of the composed functor $H_{I(Z)}^0 \circ \text{Hom}_R(N, -)$, cf. [20, Exposé VI] for details, we have in degree 0 an exact sequence

$$\begin{aligned} 0 \rightarrow {}_0\text{Hom}_{I(Z)}(I/I^2, I/I^2) &\rightarrow {}_0\text{Hom}_A(I/I^2, I/I^2) \\ &\rightarrow H^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{J}_X/\mathcal{J}_X^2, \mathcal{J}_X/\mathcal{J}_X^2)) \rightarrow {}_0\text{Ext}_{I(Z)}^1(I/I^2, I/I^2) \rightarrow \end{aligned}$$

where ${}_0\text{Hom}_{I(Z)}(I/I^2, I/I^2) \cong {}_0\text{Hom}(I/I^2, H_{I(Z)}^0(I/I^2))$ and the terms in

$$0 \rightarrow {}_0\text{Ext}^1(I/I^2, H_{I(Z)}^0(I/I^2)) \rightarrow {}_0\text{Ext}_{I(Z)}^1(I/I^2, I/I^2) \rightarrow {}_0\text{Hom}(I/I^2, H_{I(Z)}^1(I/I^2))$$

vanish by the assumptions (4.2). Hence, we conclude

$$H^0(X \setminus Z, \mathcal{H}om_{\mathcal{O}_X}(\mathcal{J}_X/\mathcal{J}_X^2, \mathcal{J}_X/\mathcal{J}_X^2)) \cong {}_0\text{Hom}_A(I/I^2, I/I^2)$$

which proves that in Theorem 4.3 we can replace the hypothesis $n \geq 2c$ by the assumptions (4.2).

Remark 4.5. If X is a linear standard determinantal scheme and $t > 1$, then $n_{1i} = t$ for all i , $n_{2j} = t + 1$ for all j , and the hypothesis $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$ is satisfied.

Remark 4.6. It is worthwhile to point out that in Theorem 4.3 the hypothesis

$$\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$$

cannot be dropped when $c = 2$. To prove it we will compute the cokernel of the morphism

$$K = R_0 \cong {}_0\text{Hom}_R(I, I) \xrightarrow{\phi_0} {}_0\text{Hom}_A(I/I^2, I/I^2).$$

To this end, we consider the diagram

$$\begin{array}{ccccc} & & & & \text{Hom}_R(I, I) \\ & & & & \downarrow \phi \\ & & & & \text{Hom}_R(I, I/I^2) \\ & & & & \downarrow \\ K_A(n+1) \otimes_R I^2 & \xleftarrow{\cong} & \text{Ext}_R^1(I, R) \otimes I^2 & \xleftarrow{\cong} & \text{Ext}_R^1(I, I^2) \\ & & \downarrow & & \downarrow \\ & & K_A(n+1) \otimes_R I & \xleftarrow{\cong} & \text{Ext}_R^1(I, I) \\ & & \downarrow \cong & & \downarrow \\ & & K_A(n+1) \otimes_R I/I^2 & \xleftarrow{\cong} & \text{Ext}_R^1(I, I/I^2) \end{array}$$

where we have used $\text{Ext}_R^2(I, -) = 0$ to get the isomorphisms. Therefore,

$$\text{coker}(\phi) = K_A(n+1) \otimes_R I^2 \cong K_A(n+1) \otimes_A I^2/I^3.$$

On the other hand for a standard determinantal scheme $X \subset \mathbb{P}^n$ of codimension 2, we have the following well-known exact sequences:

$$0 \rightarrow G^* = \bigoplus_{j=1}^t R(-n_{2j}) \rightarrow F^* = \bigoplus_{i=1}^{t+1} R(-n_{1i}) \rightarrow I \rightarrow 0$$

and

$$(4.4) \quad 0 \rightarrow \bigwedge^2 G^* \rightarrow G^* \otimes F^* \rightarrow S_2 F^* \rightarrow S_2 I \rightarrow 0.$$

Since a generic complete intersection of codimension 2 is syzygetic, i.e. $S_2 I \cong I^2$, the exact sequence

$$\cdots \rightarrow F \rightarrow G \rightarrow K_A(n+1) \rightarrow 0$$

leads to a commutative diagram

$$\begin{array}{ccccccc} & & & & G \otimes_R G^* \otimes_R F^* & & \\ & & & & \downarrow & & \\ F \otimes_R S_2 F^* & \longrightarrow & G \otimes_R S_2 F^* & & & & \\ \downarrow & & \downarrow & & & & \\ F \otimes_R I^2 & \longrightarrow & G \otimes_R I^2 & \longrightarrow & K_A(n+1) \otimes_R I^2 & \longrightarrow & 0. \end{array}$$

It follows that

$$\begin{aligned} (F \otimes S_2 F^*) \oplus (G \otimes G^* \otimes F^*) &\rightarrow G \otimes S_2 F^* = \bigoplus_{\substack{1 \leq i \leq j \leq t+1 \\ 1 \leq k \leq t}} R(-n_{1i} - n_{1j} + n_{2k}) \\ &\rightarrow K_A(n+1) \otimes_R I^2 \rightarrow 0 \end{aligned}$$

is exact and ϕ_0 is not surjective e.g. in the case $(n_{21} = n_{22} = 4$ and $n_{11} = 2 < n_{12} = n_{13} = 3)$

$$0 \rightarrow R(-4)^2 \rightarrow R(-3)^2 \oplus R(-2) \rightarrow I \rightarrow 0.$$

We have stated Theorem 4.3 for standard determinantal schemes because the paper concerns the main features of the normal sheaf of a standard determinantal scheme $X \subset \mathbb{P}^n$. Nevertheless, the result works in a much more general set up. Indeed, the assumption that X is a standard determinantal scheme is not necessary and the result and its proof hold for any graded quotient of the polynomial ring provided $\text{depth}_{I(Z)} I/I^2 \geq 2$. In fact, it holds

Theorem 4.7. *Let $X \subset \mathbb{P}^n$ be a closed subscheme of codimension $c \geq 2$ (not necessarily ACM) with a minimal free R -resolution*

$$\cdots \rightarrow \bigoplus_j^{b_2} R(-n_{2j}) \rightarrow \bigoplus_i^{b_1} R(-n_{1i}) \rightarrow R \rightarrow R/I \rightarrow 0$$

where $I := I(X)$. Let $Z \subset X$ be a closed subset such that $X \setminus Z \hookrightarrow \mathbb{P}^n$ is a local complete intersection. Let L be a finitely generated R/I -module that is invertible over $X \setminus Z$, put

$$\mathcal{N}_X(\mathcal{L}) := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{L})$$

and assume $\text{depth}_{I(Z)} L \geq 2$, $\text{depth}_{I(Z)} I/I^2 \geq 2$ and $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$. Then,

$$\text{Hom}(\mathcal{N}_X(\mathcal{L}), \mathcal{N}_X(\mathcal{L})) \cong {}_0\text{Hom}_A(I/I^2, I/I^2) \cong K.$$

Remark 4.8. For a complete intersection of codimension $c \geq 2$ and dimension $n-c \geq 1$, the conclusion is false while all assumptions, except for $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$ are obviously satisfied.

Remark 4.9. As explained in Remark 4.4, the hypothesis $\text{depth}_{I(Z)} I/I^2 \geq 2$ in Theorem 4.7 can be replaced by

$$(4.5) \quad H_{I(Z)}^1(I/I^2)_{n_{1i}} = H_{I(Z)}^0(I/I^2)_{n_{1i}} = H_{I(Z)}^0(I/I^2)_{n_{2j}} = 0 \quad \text{for any } i \text{ and } j.$$

As an application we have:

Corollary 4.10. *Let $X \subset \mathbb{P}^n$ be either a codimension 2 ACM subscheme with a minimal free R -resolution*

$$0 \rightarrow \bigoplus_j^v R(-n_{2j}) \rightarrow \bigoplus_i^{v+1} R(-n_{1i}) \rightarrow R \rightarrow R/I \rightarrow 0$$

or a codimension 3 arithmetically Gorenstein subscheme with a minimal free R -resolution

$$0 \rightarrow R(-e) \rightarrow \bigoplus_j^v R(-n_{2j}) \rightarrow \bigoplus_i^v R(-n_{1i}) \rightarrow R \rightarrow R/I \rightarrow 0.$$

Let $Z \subset X$ be a closed subset such that $X \setminus Z \hookrightarrow \mathbb{P}^n$ is a local complete intersection. Assume

$$\text{depth}_{I(Z)} R/I \geq \begin{cases} 3 & \text{if } c = 2 \\ 2 & \text{if } c = 3 \end{cases} \quad \text{and} \quad \max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}.$$

Then, \mathcal{N}_X is simple.

Proof. By Theorem 4.7 it suffices to show that $\text{depth}_{I(Z)} I/I^2 \geq 2$. For $c = 2$, we get

$$\text{depth}_{I(Z)} I/I^2 \geq \text{depth}_{I(Z)} R/I - 1$$

by [1], cf. (4.4). For $c = 3$ we know that I/I^2 is MCM by [7] because $I/I^2 \otimes K_A$ is MCM in the licci case and the canonical module K_A is trivial in the Gorenstein case. So, we are done \square

Let us give some examples of ACM schemes $X \subset \mathbb{P}^N$ with simple normal sheaf.

Example 4.11. Let $X \subset \mathbb{P}^3$ be a (smooth) ACM curve. Since

$$\text{depth } I/I^2 \geq \dim R/I - 1,$$

we have $H_{\mathfrak{m}}^0(I/I^2)_{\mu} = 0$ for any μ . So, according to Theorem 4.3 and Remark 4.4, we only need to check

$$(4.6) \quad H_{\mathfrak{m}}^1(I/I^2)_{n_{1i}} \cong H^1(X, \mathcal{I}_X^2(n_{1i})) = 0 \quad \text{for any } i$$

and

$$\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$$

to conclude that \mathcal{N}_X is simple. Using Macaulay2, we get

- (i) (4.6) does not hold in the linear case.
- (ii) If $\deg \mathcal{A} = \begin{pmatrix} 1 & 1 & 2 \\ 1 & 1 & 2 \end{pmatrix}$, then (4.6) holds. However, since the condition

$$4 = \max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\} = 4$$

is not true, we cannot conclude that \mathcal{N}_X is simple. In fact, we have seen in Remark 4.6 that \mathcal{N}_X is not simple.

- (iii) If $\deg \mathcal{A} = \begin{pmatrix} 1 & 2 & 2 \\ 1 & 2 & 2 \end{pmatrix}$ or $\begin{pmatrix} 2 & 2 & 2 \\ 2 & 2 & 2 \end{pmatrix}$ or $\begin{pmatrix} 3 & 2 & 2 \\ 3 & 2 & 2 \end{pmatrix}$ or $\begin{pmatrix} 3 & 3 & 1 \\ 3 & 3 & 1 \end{pmatrix}$, then (4.6) holds as well as the inequality $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$; and we get that \mathcal{N}_X is simple.

In conclusion, the assumptions (4.6) and $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$ seem weak for ACM curves in \mathbb{P}^3 .

Example 4.12. Let $X \subset \mathbb{P}^4$ be a smooth standard determinantal curve. By Theorem 4.3 and Remark 4.4, to prove that \mathcal{N}_X is simple, we only need to check that the following hypothesis are satisfied:

$$(4.7) \quad H_{\mathfrak{m}}^1(I/I^2)_{n_{1i}} = H_{\mathfrak{m}}^0(I/I^2)_{n_{1i}} = H_{\mathfrak{m}}^0(I/I^2)_{n_{2j}} = 0 \quad \text{for any } i \text{ and } j$$

and

$$\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}.$$

Using Macaulay2, we get

- (i) (4.7) does not hold in the linear case.
- (ii) If $\deg \mathcal{A} = \left(1 \frac{2}{1} \frac{2}{2} \frac{2}{2}\right)$, then (4.7) holds while $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$ is not true and we cannot conclude that \mathcal{N}_X is simple.
- (iii) If $\deg \mathcal{A} = \left(\frac{2}{2} \frac{2}{2} \frac{2}{2} \frac{2}{2}\right)$, then (4.7) holds as well as the inequality $\max\{n_{2j}\} < 2 \cdot \min\{n_{1i}\}$. Hence, \mathcal{N}_X is simple.

We will end this section with a result about the indecomposability of the normal sheaf of a standard determinantal scheme $X \subset \mathbb{P}^n$ which does not involve the degrees of the generators (resp. first syzygies) of $I(X)$. To achieve our goal we need the following preliminary lemma.

Lemma 4.13. *Let $X \subset \mathbb{P}^n$ be a codimension c standard determinantal subscheme associated to a graded morphism $\varphi : F \rightarrow G$. Set $M = \text{coker}(\varphi)$ and $J = I_{t-1}(\varphi)$. Let*

$$H \cong \mathbb{P}^{n-1} \subset \mathbb{P}^n$$

be a hyperplane defined by a general form $h \in R_1$. Consider $X' = X \cap H \subset H$ the hyperplane section of X and $R' = R/(h) \cong K[z_0, \dots, z_{n-1}]$. Then X' is a codimension c standard determinantal subscheme of \mathbb{P}^{n-1} associated to the graded morphism

$$\varphi' = \varphi \otimes 1 : F' := F \otimes_R R/(h) \rightarrow G' := G \otimes_R R/(h)$$

and $M' := \text{coker}(\varphi') \cong M \otimes_R R/(h)$. Assume $\text{depth}_J A \geq 3$. Then, we have

$$\text{Ext}_R^1(M, S_i M) \otimes_R R/(h) \cong \text{Ext}_{R'}^1(M', S_i M') \quad \text{for } 1 \leq i \leq c.$$

Proof. Let us consider the free R -resolution W_\bullet of $\text{Ext}_R^1(M, S_i M)$ of Corollary 3.7,

$$\begin{aligned} 0 \rightarrow Q_c \rightarrow Q_{c-1} \oplus P_c \rightarrow Q_{c-2} \oplus P_{c-1} \oplus F_c \rightarrow Q_{c-3} \oplus P_{c-2} \oplus F_{c-1} \\ \rightarrow \cdots \rightarrow Q_0 \oplus P_1 \oplus F_2 \rightarrow P_0 \oplus F_1 \rightarrow F_0 \rightarrow \text{Ext}_R^1(M, S_i M) \rightarrow 0, \end{aligned}$$

and the free R' -resolution W'_\bullet of $\text{Ext}_{R'}^1(M', S_i M')$,

$$\begin{aligned} 0 \rightarrow Q'_c \rightarrow Q'_{c-1} \oplus P'_c \rightarrow Q'_{c-2} \oplus P'_{c-1} \oplus F'_c \rightarrow Q'_{c-3} \oplus P'_{c-2} \oplus F'_{c-1} \\ \rightarrow \cdots \rightarrow Q'_0 \oplus P'_1 \oplus F'_2 \rightarrow P'_0 \oplus F'_1 \rightarrow F'_0 \rightarrow \text{Ext}_{R'}^1(M', S_i M') \rightarrow 0, \end{aligned}$$

also given by Corollary 3.7 because $\text{depth}_J A \geq 3$. Since $h \in R_1$ is a general linear form and $\text{depth}_m \text{Ext}_R^1(M, S_i M) \geq 1$ by Corollary 3.7, we have

$$\text{Ext}_R^1(M, S_i M) : h = \text{Ext}_R^1(M, S_i M)$$

and by [5, Lemma 1.3.5] that

$$W_\bullet \otimes_R R/(h) = W'_\bullet$$

is a free $R' = R/(h)$ -resolution of $\text{Ext}_R^1(M, S_i M) \otimes_R R/(h)$. Therefore, we conclude that

$$\text{Ext}_R^1(M, S_i M) \otimes_R R/(h) \cong \text{Ext}_{R'}^1(M', S_i M'). \quad \square$$

Now, we are ready to prove the indecomposability of the normal sheaf of a standard determinantal scheme $X \subset \mathbb{P}^n$ under some mild hypothesis which does not involve the degrees

of the generators (resp. first syzygies) of $I(X)$. Recalling

$$\mathcal{N}_X(\mathcal{M}^i) := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}_X/\mathcal{I}_X^2, \widetilde{S}_i M),$$

we have

Theorem 4.14. *Let $X \subset \mathbb{P}^n$ be a standard determinantal scheme of codimension $c \geq 2$. We keep the notation introduced above and set $J = I_{t-1}(\mathcal{A})$. Assume $a_j - b_i > 0$ for all i, j and*

$$\text{depth}_J A \geq \begin{cases} c + 2 & \text{if } c > 2, \\ 3 & \text{if } c = 2. \end{cases}$$

If $c = 2$ and $n \geq 4$ or $c \geq 3$ and $n \geq 2c + 1$, then the normal sheaf \mathcal{N}_X and more generally, the “twisted” normal sheaves $\mathcal{N}_X(\mathcal{M}^i)$, $-1 \leq i \leq c - 1$, are indecomposable.

Proof. The idea is to fix c and use induction on n . The initial cases are true. In fact, if $(c, n) = (2, 4)$, then X is smooth because $\text{depth}_J A = \text{depth}_{\mathfrak{m}} A$ and \mathcal{N}_X is indecomposable by [3, Théorème A]. If we have $(c, n) = (c, 2c + 1)$, $c \geq 3$, then X is smooth, $\text{Pic}(X) \cong \mathbb{Z}$ (cf. Theorem 2.4) and \mathcal{N}_X is indecomposable by [2, Theorem 3.2]. It follows that

$$\mathcal{N}_X(\mathcal{M}^i) \cong (\mathcal{I}_X/\mathcal{I}_X^2)^\vee \otimes \widetilde{S}_i M$$

is indecomposable. The result now follows from induction using Lemma 4.13 and taking into account that

$$\text{Ext}_R^1(M, S_{i+1}M) \cong \text{Hom}_A(I/I^2, S_i M)$$

(see Proposition 3.5 and Remark 3.10). □

Example 4.15. (1) We consider $X \subset \mathbb{P}^n$, $n \geq 4$, the standard determinantal subscheme of codimension 2 associated to the matrix

$$\begin{pmatrix} x_0 & x_1 & x_2^2 \\ x_3 & x_4 & f \end{pmatrix}$$

where f is a general form of degree 2. We have seen in Remark 4.6 that \mathcal{N}_X is not simple but it follows from Theorem 4.14 that \mathcal{N}_X is indecomposable.

(2) We consider a rational normal scroll $S(a_0, \dots, a_k)$, i.e. the image of the map

$$\sigma : \mathbb{P}^1 \times \mathbb{P}^k \rightarrow \mathbb{P}^N$$

given by

$$\sigma(x, y; t_0, t_1, \dots, t_k) := (x^{a_0}t_0, x^{a_0-1}yt_0, \dots, y^{a_0}t_0, \dots, x^{a_k}t_k, x^{a_k-1}yt_k, \dots, y^{a_k}t_k)$$

where $N = k + \sum_{i=0}^k a_i$. If we choose coordinates $X_0^0, \dots, X_{a_0}^0, \dots, X_0^k, \dots, X_{a_k}^k$ in \mathbb{P}^N , the ideal of $S(a_0, \dots, a_k)$ is generated by the maximal minors of the $2 \times c$ matrix with two rows and $k + 1$ catalecticant blocks:

$$M_{a_0, \dots, a_k} := \begin{pmatrix} X_0^0 & \cdots & X_{a_0-1}^0 & \cdots & X_0^k & \cdots & X_{a_k-1}^k \\ X_1^0 & \cdots & X_{a_0}^0 & \cdots & X_1^k & \cdots & X_{a_k}^k \end{pmatrix}.$$

By Theorem 4.14, the normal bundle of $S(1, \dots, 1)$ and of $S(2, 1, \dots, 1)$ are indecomposable. (Notice that $S(1, \dots, 1)$ corresponds to the Segre variety $\mathbb{P}^1 \times \mathbb{P}^k \hookrightarrow \mathbb{P}^{2k+1}$ already discussed in [2, Corollary 3.3].)

Arguing as in the proof of Theorem 4.14, we get:

Corollary 4.16. *Let $X \subset \mathbb{P}^n$ be a smooth standard determinantal scheme of codimension c and let $\mathbb{P}^{n-1} \cong H \subset \mathbb{P}^n$ a general hyperplane and set $X' = X \cap H$. Assume $n - c \geq 2$ and $\text{depth}_J A \geq 3$, $J = I_{t-1}(A)$. Then, $(\mathcal{N}_{X/\mathbb{P}^n})|_{\mathbb{P}^{n-1}} \cong \mathcal{N}_{X'/\mathbb{P}^{n-1}}$.*

5. The μ -semistability of the normal sheaf of a determinantal varieties

The normal bundle of a smooth variety $X \subset \mathbb{P}^n$ has been intensively studied since it reflects many properties of the embedding; so far few examples of smooth varieties having μ -(semi)stable normal bundle are known. The first example of a curve $C \subset \mathbb{P}^3$ with normal bundle \mathcal{N}_C μ -stable was given by Sacchiero ([35]). In [16, Proposition 2], Ellia proved that the normal bundle of a linear determinantal curve $C \subset \mathbb{P}^3$ is linear μ -semistable and the normal bundle \mathcal{N}_C of a general ACM curve $C \subset \mathbb{P}^3$ of degree 6 and genus 3 is μ -stable (see [17, 34] for more information about the μ -stability of the normal sheaf of a curve in \mathbb{P}^3). The first goal of this last section is to generalize Ellia's result for linear determinantal curves C in \mathbb{P}^3 to linear determinantal schemes $X \subset \mathbb{P}^n$ of arbitrary dimension.

As in previous sections we will assume $c > 1$, since the case $c = 1$ corresponds to a hypersurface $X \subset \mathbb{P}^n$ and $\mathcal{N}_X \cong \mathcal{O}_X(\delta)$, $\delta := \deg(X)$, is μ -stable. We will also assume $t > 1$, since the case $t = 1$ corresponds to a codimension c complete intersection $X \subset \mathbb{P}^n$ and $\mathcal{N}_X = \bigoplus_{i=1}^c \mathcal{O}_X(d_i)$, $d_i \in \mathbb{Z}$, is not μ -stable.

Let us start recalling the definition of μ -(semi)stability

Definition 5.1. Let $X \subset \mathbb{P}^n$ be a smooth projective scheme of dimension d and let \mathcal{E} be a coherent sheaf on X . \mathcal{E} is said to be μ -semistable if for any non-zero coherent subsheaf \mathcal{F} of \mathcal{E} we have the inequality

$$\mu(\mathcal{F}) := \frac{\deg(c_1(\mathcal{F}))}{\text{rk}(\mathcal{F})} \leq \mu(\mathcal{E}) := \frac{\deg(c_1(\mathcal{E}))}{\text{rk}(\mathcal{E})}$$

where as usual $\deg(c_1(\mathcal{F})) = c_1(F) \cdot H^{d-1}$. We say that \mathcal{E} is μ -stable if strict inequality “ $<$ ” always holds.

Remark 5.2. Recall that μ -stable sheaves are simple and hence indecomposable but not vice versa.

Theorem 5.3. *Let $X \subset \mathbb{P}^n$ be a smooth linear determinantal scheme of codimension $c \geq 2$ associated to a $t \times (t + c - 1)$ matrix A . Assume $n - c \geq 1$. Then, \mathcal{N}_X is μ -semistable.*

Proof. Since the notion of μ -semistability is preserved when we twist by an invertible sheaf, it will be enough to prove that $\mathcal{N}_X(-H) \otimes \widetilde{S_{c-2}M}$ is μ -semistable. It follows from Theorem 3.6 that $\mathcal{N}_X(-H) \otimes \widetilde{S_{c-2}M}$ is a rank c Ulrich sheaf on X and by [9, Theorem 2.9] any Ulrich sheaf is μ -semistable which proves what we want. \square

Remark 5.4. Since Ulrich sheaves are Gieseker semistable, the above proof also shows that the normal sheaf \mathcal{N}_X to a smooth linear determinantal scheme of codimension $c \geq 2$ is Gieseker semistable.

Remark 5.5. Without extra hypothesis the above result cannot be improved and the μ -stability of the normal sheaf \mathcal{N}_X of a linear standard determinantal scheme $X \subset \mathbb{P}^n$ cannot be guaranteed. In fact, if we consider a rational normal curve $C \subset \mathbb{P}^n$ defined by the 2×2 minors of a $2 \times n$ matrix with general linear entries, it is well known that $\mathcal{N}_{X/\mathbb{P}^n} \cong \mathcal{O}_X(2)^{n-1}$. Therefore, \mathcal{N}_X is μ -semistable but not μ -stable.

In the codimension 2 case, Theorem 5.3 can be improved using the following lemma:

Lemma 5.6. *Let $X \subset \mathbb{P}^n$ be a smooth linear determinantal scheme of codimension $c \geq 2$ and dimension $n - c \geq 2$ defined by the maximal minors of a $t \times (t + c - 1)$ matrix \mathcal{A} . Assume $t \geq n$ when $n - c = 2$. Let H be a general hyperplane section of X and let $Y \subset \mathbb{P}^n$ be the codimension 1 subscheme of X defined by the maximal minors of the $t \times (t + c)$ matrix \mathcal{B} obtained adding to \mathcal{A} a column of general linear forms. Let \mathcal{L} be a line bundle on X . It holds:*

- (i) \mathcal{L} is an ACM line bundle on X if and only if $\mathcal{L} \cong \mathcal{O}_X(aY + bH)$ with $-1 \leq a \leq c$ and $b \in \mathbb{Z}$,
- (ii) \mathcal{L} is an initialized Ulrich line bundle if and only if $\mathcal{L} \cong \mathcal{O}_X(-Y + tH)$ or $\mathcal{O}_X(cY - cH)$.

Proof. See [27]. □

In fact, we have:

Theorem 5.7. *Let X be a smooth linear determinantal scheme of codimension 2 in \mathbb{P}^n defined by the maximal minors of a $t \times (t + 1)$ matrix with linear entries. Assume that $n \geq 4$. Then, \mathcal{N}_X is μ -stable. In particular, \mathcal{N}_X is simple and indecomposable.*

Proof. For the case $(n, t) = (4, 3)$ the reader can see [32, Proposition 4.10]. Assume that $(n, t) \neq (4, 3)$. Since μ -(semi)stability is preserved when we twist by line bundles, we know that $\mathcal{N}_X(-H)$ is μ -semistable (Theorem 5.3) and we want to prove that it is μ -stable, i.e. we must rule out the existence of a coherent subsheaf $\mathcal{F} \subset \mathcal{N}_X(-H)$ with the property that $\mu(\mathcal{F}) = \mu(\mathcal{N}_X(-H))$. By pulling-back torsion, if necessary, we may assume that $\mathcal{N}_X(-H)/\mathcal{F}$ is torsion free in which case \mathcal{F} is locally free and we have an exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{N}_X(-H) \rightarrow \mathcal{N}_X(-H)/\mathcal{F} \rightarrow 0$$

of coherent sheaves with $\mathcal{N}_X(-H)/\mathcal{F}$ torsion free and $\mu(\mathcal{F}) = \mu(\mathcal{N}_X(-H))$. By [9, Theorem 2.9 (b)], \mathcal{F} and $\mathcal{N}_X(-H)/\mathcal{F}$ are both Ulrich line bundles. So, according to our Lemma 5.6 we have four possibilities:

- (1) $0 \rightarrow \mathcal{O}_X(-Y + tH) \rightarrow \mathcal{N}_X(-H) \rightarrow \mathcal{O}_X(-Y + tH) \rightarrow 0$,
- (2) $0 \rightarrow \mathcal{O}_X(2Y - 2H) \rightarrow \mathcal{N}_X(-H) \rightarrow \mathcal{O}_X(-Y + tH) \rightarrow 0$,
- (3) $0 \rightarrow \mathcal{O}_X(-Y + tH) \rightarrow \mathcal{N}_X(-H) \rightarrow \mathcal{O}_X(2Y - 2H) \rightarrow 0$,
- (4) $0 \rightarrow \mathcal{O}_X(2Y - 2H) \rightarrow \mathcal{N}_X(-H) \rightarrow \mathcal{O}_X(2Y - 2H) \rightarrow 0$.

Let us check that none of them is allowed. To this end, we will start computing the Chern classes of $\mathcal{N}_X(-H)$. We sheafify the exact sequence (3.3) and we get the exact sequence

$$(5.1) \quad 0 \rightarrow \mathcal{O}_X(-H) \rightarrow \mathcal{O}_X(Y - 2H)^t \rightarrow \mathcal{O}_X(Y - H)^{t+1} \rightarrow \mathcal{N}_X(-H) \rightarrow 0.$$

Therefore, the Chern polynomial $c_u(\mathcal{N}_X(-H))$ of $\mathcal{N}_X(-H)$ is given by

$$\begin{aligned} c_u(\mathcal{N}_X(-H)) &= \sum_u c_i(\mathcal{N}_X(-H))u^i \\ &= \frac{(1 - Hu)(1 + (Y - H)u)^{t+1}}{(1 + (Y - 2H)u)^t} \\ &= \frac{(1 - Hu)(\sum_{k=0}^{t+1} \binom{t+1}{k}(Y - H)^k u^k)}{\sum_{k=0}^t \binom{t}{k}(Y - 2H)^k u^k} \end{aligned}$$

and a straightforward computation gives us

$$c_1(\mathcal{N}_X(-H)) = Y + (t - 2)H$$

and

$$c_2(\mathcal{N}_X(-H)) = -YH + \frac{t^2 - t + 2}{2}H^2.$$

Comparing the first Chern class we eliminate the possibilities (1) and (4) because in case (1) we would get $c_1(\mathcal{N}_X(-H)) = -2Y + 2tH$ and in case (4) we would get $c_1(\mathcal{N}_X(-H)) = 4Y - 4H$. To rule out the two remaining cases, we compare the second Chern class; in both cases we would get $c_2(\mathcal{N}_X(-H)) = 2(1 + t)YH - 2Y^2 - 2tH^2$ which is impossible and this concludes the proof of the theorem. \square

References

- [1] *L. Avramov and J. Herzog*, The Koszul algebra of a codimension 2 embedding, *Math. Z.* **175** (1980), 249–260.
- [2] *L. Badescu*, On the normal bundle of submanifolds of \mathbb{P}^2 , *Proc. Amer. Math. Soc.* **136** (2008), 1505–1513.
- [3] *B. Basili and C. Peskine*, Décomposition du fibré normal des surfaces lisses de \mathbb{P}^4 et structures doubles sur les solides de \mathbb{P}^5 , *Duke Math. J.* **69** (1993), 87–95.
- [4] *J. Brennan, J. Herzog and B. Ulrich*, Maximally generated Cohen–Macaulay modules, *Math. Scand.* **61** (1987), 181–203.
- [5] *W. Bruns and J. Herzog*, Cohen–Macaulay rings, *Cambridge Stud. Adv. Math.* **39**, Cambridge University Press, Cambridge 1993.
- [6] *W. Bruns and U. Vetter*, Determinantal rings, *Lectures Notes in Math.* **1327**, Springer-Verlag, New York 1988.
- [7] *R. O. Buchweitz*, Contributions à la théorie des singularités, Thesis l’Université Paris VII, 1981.
- [8] *M. Casanellas and R. Hartshorne*, Gorenstein biliaison and ACM sheaves, *J. Algebra* **278** (2004), 314–341.
- [9] *M. Casanellas and R. Hartshorne*, Stable Ulrich bundles, preprint 2011, <http://arxiv.org/abs/1102.0878>.
- [10] *L. Costa, R. M. Miró-Roig and J. Pons-Llopis*, The representation type of Segre varieties, *Adv. Math.* **230** (2012), 1995–2013.
- [11] *Y. Drozd and G. M. Greuel*, Tame and wild projective curves and classification of vector bundles, *J. Algebra* **246** (2001), 1–54.
- [12] *L. Ein*, On the cohomology of projective Cohen–Macaulay determinantal varieties of \mathbb{P}^n , in: *Geometry of complex projective varieties (Cetraro 1990)*, *Semin. Conf.* **9**, Mediterranean Press, Rende (1993), 143–152.
- [13] *D. Eisenbud*, *Commutative algebra. With a view toward algebraic geometry*, *Grad. Texts in Math.* **150**, Springer-Verlag, New York 1995.

- [14] *D. Eisenbud* and *F. Schreyer*, Boij–Söderberg theory, in: Combinatorial aspects of commutative algebra and algebraic geometry. The Abel symposium 2009 (Voss 2009), Abel Symp. **6**, Springer-Verlag, Berlin (2011), 35–48
- [15] *D. Eisenbud*, *F. Schreyer* and *J. Weyman*, Resultants and chow forms via exterior syzygies, J. Amer. Math. Soc. **16** (2003), 537–579.
- [16] *P. Ellia*, Exemples de courbes de \mathbb{P}^3 à fibré normal semi-stable, stable Math. Ann. **264** (1983), no. 3, 389–396.
- [17] *G. Ellingsrud* and *A. Hirschowitz*, Sur le fibré normal des courbes gauches, C.R. Acad. Sci. Paris Sér. I Math. **299** (1984), 245–248.
- [18] *D. Faenzi* and *M. L. Fania*, On the Hilbert scheme of determinantal subvarieties, preprint 2010, <http://arxiv.org/abs/1012.4692>; to appear in Math. Res. Lett.
- [19] *D. Grayson* and *M. Stillman*, Macaulay 2—A software system for algebraic geometry and commutative algebra, <http://www.math.uiuc.edu/Macaulay2/>.
- [20] *A. Grothendieck*, Cohomologie locale des faisceaux cohérents et Théorèmes de Lefschetz locaux et globaux, North Holland, Amsterdam 1968.
- [21] *G. Horrocks*, Vector bundles on the punctured spectrum of a local ring, Proc. Lond. Math. Soc. (3) **14** (1964), 689–713.
- [22] *D. Kirby*, A sequence of complexes associated to a matrix, J. Lond. Math. Soc. (2) **7** (1973), 523–530.
- [23] *J. O. Kleppe*, Unobstructedness and dimension of families of Gorenstein algebras, Collect. Math. **58** (2007), 199–238.
- [24] *J. O. Kleppe*, Deformations of modules of the maximal grade and Hilbert scheme at determinantal schemes, J. Algebra **407** (2014), 246–274.
- [25] *J. O. Kleppe* and *R. M. Miró-Roig*, Dimension of families of determinantal schemes, Trans. Amer. Math. Soc. **357** (2005), 2871–2907.
- [26] *J. O. Kleppe* and *R. M. Miró-Roig*, Families of determinantal schemes, Proc. Amer. Math. Soc. **139** (2011), 3831–3843.
- [27] *J. O. Kleppe* and *R. M. Miró-Roig*, in preparation.
- [28] *J. O. Kleppe* and *C. Peterson*, Maximal Cohen–Macaulay modules and Gorenstein algebras, J. Algebra **238** (2001), 776–800.
- [29] *J. O. Kleppe* and *C. Peterson*, Sheaves with canonical determinant on Cohen–Macaulay schemes, J. Algebra **256** (2002), 250–279.
- [30] *A. López*, Noether–Lefschetz theory and the Picard group of projective surfaces, Mem. Amer. Math. Soc. **438** (1991).
- [31] *R. M. Miró-Roig*, The representation type of rational normal scrolls, Rend. Circ. Mat. Palermo (2) **62** (2013), 153–164.
- [32] *R. M. Miró-Roig* and *J. Pons-Llopis*, Representation type of rational ACM surfaces $X \subseteq \mathbb{P}^4$, Algebr. Represent. Theory **16** (2013), 1135–1157.
- [33] *R. M. Miró-Roig* and *J. Pons-Llopis*, n -dimensional Fano varieties of wild representation type, J. Pure Appl. Algebra **218** (2014), no. 10, 1867–1884.
- [34] *D. Perrin*, Courbes passant par m points généraux de \mathbb{P}^3 , Mém. Soc. Math. France **28–29** (1987).
- [35] *G. Sacchiero*, Exemple de courbes de \mathbb{P}^3 de fibré normal stable, Comm. Algebra **11** (1983), 2115–2121.
- [36] *B. Ulrich*, Gorenstein rings and modules with high number of generators, Math. Z. **188** (1984), 23–32.

Jan O. Kleppe, Faculty of Technology, Art and Design, Oslo and Akershus University College,
PB. 4, St. Olavs Plass, N-0130 Oslo, Norway
e-mail: JanOddvar.Kleppe@hioa.no

Rosa M. Miró-Roig, Departament d’Algebra i Geometria, Facultat de Matemàtiques,
Gran Via de les Corts Catalanes 585, 08007 Barcelona, Spain
e-mail: miro@ub.edu

Eingegangen 30. Juli 2013, in revidierter Fassung 12. März 2014