

STUDY PROTOCOL

Open Access



A falls prevention programme to improve quality of life, physical function and falls efficacy in older people receiving home help services: study protocol for a randomised controlled trial

Maria Bjerk^{1*} , Therese Brovold¹, Dawn A. Skelton² and Astrid Bergland¹

Abstract

Background: Falls and fall-related injuries in older adults are associated with great burdens, both for the individuals, the health care system and the society. Previous research has shown evidence for the efficiency of exercise as falls prevention. An understudied group are older adults receiving home help services, and the effect of a falls prevention programme on health-related quality of life is unclear. The primary aim of this randomised controlled trial is to examine the effect of a falls prevention programme on quality of life, physical function and falls efficacy in older adults receiving home help services. A secondary aim is to explore the mediating factors between falls prevention and health-related quality of life.

Methods: The study is a single-blinded randomised controlled trial. Participants are older adults, aged 67 or older, receiving home help services, who are able to walk with or without walking aids, who have experienced at least one fall during the last 12 months and who have a Mini Mental State Examination of 23 or above. The intervention group receives a programme, based on the Otago Exercise Programme, lasting 12 weeks including home visits and motivational telephone calls. The control group receives usual care. The primary outcome is health-related quality of life (SF-36). Secondary outcomes are leg strength, balance, walking speed, walking habits, activities of daily living, nutritional status and falls efficacy. All measurements are performed at baseline, following intervention at 3 months and at 6 months' follow-up. Sample size, based on the primary outcome, is set to 150 participants randomised into the two arms, including an estimated 15–20% drop out. Participants are recruited from six municipalities in Norway.

Discussion: This trial will generate new knowledge on the effects of an exercise falls prevention programme among older fallers receiving home help services. This knowledge will be useful for clinicians, for health managers in the primary health care service and for policy makers.

Trial registration: ClinicalTrials.gov. NCT02374307. First registration, 16/02/2015.

Keywords: Falls prevention, Home help services, Elderly, Quality of life, Older adults, Exercise, Balance, Preventative care

* Correspondence: maria.bjerk@hioa.no

¹Department of Physiotherapy, Faculty of Health Sciences, Oslo and Akershus University College, PO box 4 St. Olavs plass, Oslo 0130, Norway
Full list of author information is available at the end of the article



Background

Older adults and health-related quality of life

Health-related quality of life (HRQOL) is of great interest, both with respect to individuals themselves as well as a primary concern of public health administrations and professionals. The remarkable increase in life expectancy in the twentieth century implies a need to focus on factors capable of promoting a high level of HRQOL into old age. In fact, older adults seem to prefer a high HRQOL more than longevity, and researchers have concluded that the key challenge is to preserve a high level of HRQOL rather than increase length of life [1, 2]. HRQOL is a subjective, multidimensional concept shaped by, but not entirely dependent upon, the effects of disease and treatment [3]. The WHO Quality of life (QOL) group defines QOL as “individuals’ perception of their position in life in the context of the culture and value systems in which they live, and in relation to their goals, expectations, standards and concerns” [4]. Public health policies in many European countries are therefore primarily concerned with keeping older people living independently in the community with a good quality of life [5–7]. The raise in number of older adults implies more people with chronic diseases and a greater challenge for the health care system in finding effective and feasible interventions to reach this goal [5–8].

Aiming to enable older people to live at home as long as possible, the municipalities in Norway are responsible for providing services in the form of home help for older people [9]. Home help includes services that assist instrumental activities of daily living (iADL), such as vacuum cleaning, and personal activities of daily living (pADL), such as getting dressed, safety alarm services to provide assistance if they fall, and social support. The most important predictor of home care use seems to be dependency in IADL and ADL and cognitive impairment [10]. Home help receivers constitute a transitional group between independent community living older people, and people living in residential care facilities/nursing homes [11]. The combination of the increase of older people and the use of the so-called LEON (“lowest most efficient level of care”) principle [6], makes this a steadily growing group in society, and can be seen as an especially vulnerable group among the older population. Moreover, as economic resources are scarce, there seems to be more focus on post-acute care instead of health promotion and prevention to maintain older adults at home [12]. To date there is no evidence-based practice standard for falls-prevention in Norwegian home care services.

Older adults and falls

Falls and fall-related injuries are common in older adults and are associated with substantial economic costs that are borne by individuals, the community, and the

medical system as a whole [13, 14]. Up to 40% of all nursing home admissions have been found to relate to falls and instability [15]. Important risk factors for falling in the group of older adults are impaired balance and gait, polypharmacy and a history of falls [16]. Poor nutritional status has also been associated with an increased risk of falling, [17] and malnutrition or being in risk of malnutrition is prevalent in half of the older adults receiving home care services [18]. Common consequences of a fall are fear of falling, activity restrictions, loss of mobility and loss of independence [19]. Falling, or being at risk of falling also has a negative influence on QOL [20]. Hence, it can be argued that HRQOL is an important outcome in the assessment of falls-prevention programmes [21].

After several decades of research on interventions to reduce falls and fall risk factors, there is now strong evidence for the effectiveness and cost-efficiency of exercise in reducing the number of falls [14, 22–25]. An important, but yet understudied group when it comes to the effect of falls prevention programmes are older adults receiving home help services, and especially those who recently have experienced a fall [11]. Previous research has shown that falls and fear of falling are common in this population and are correlated with the amount of home care needed [11, 26]. Vikman et al. [11] concluded that future studies should have a focus on the effects of falls prevention programmes in the group of those receiving home help services. Recently it has been shown that home help receivers fell more frequently than the independent home-dwelling older population [27]. Low functional level and high home care recipient health problems were independently associated with risk of falling [27]. Fear of falling is also reported more frequently in the group of older adults receiving home help services compared to those who do not receive home care [28]. This suggests that the higher level of fear of falling could be due to a higher level of frailty in this group. Finally, it has been shown that elderly home help receivers in Sweden have a lower QOL compared to those without help and that QOL was negatively correlated with the amount of help needed [29].

Interventions to improve quality of life

Although exercise-based falls prevention programmes have shown a clear effect on falls incidence and fall risk factors in the general older population, the evidence is still inconsistent about the effects on HRQOL and in particular related to the population of home help receivers [11, 21]. A systematic review by Vaapio et al. [21] considered the specific effect of falls prevention programmes on QOL. The review looked at 12 RCTs including older adults, but none of the studies were aimed at home help receivers. Six of these studies showed a

positive effect on QOL. The interventions in these studies ranged from exercise (two studies), information based (one study), to comprehensive geriatric assessment (one study). The review concluded that there is a lack of evidence about the potential benefits of falls prevention programmes on QOL in older people and that more research is needed.

To the authors' knowledge, only two RCTs have been examining exercise interventions aimed specifically at the population of older home help receivers [30, 31]. The first study tested a home-exercise programme and found positive results on maximum walking speed, but unfortunately the assessors were not blinded to the intervention [31]. The other study explored the effects and costs of a multifactorial, interdisciplinary team approach to falls prevention in 109 older home help receivers with a risk for falls [30]. Exercise was part of the programme, but the amount and mode of exercise varied according to individual needs. At 6 months, no difference in the mean number of falls between groups were found. Subgroup analyses showed that the intervention effectively reduced falls in men (75–84 years old) with a fear of falling or negative fall history, but it is unclear whether the study had sufficiently power for subgroup analyses [30]. Nevertheless, the secondary outcome of QOL significantly improved in the intervention group.

The effect of exercise interventions on HRQOL in the general older adult population have had mixed results, reporting both statistically significant positive effects as well as no significant changes [32–35]. A meta-analysis found no difference between aerobic and strength training, suggesting that the different exercise modes yielded the same effect on self-reported physical function domains of HRQOL [34]. Acree et al. [3] concluded that healthy older adults who regularly participated in physical activity of at least moderate intensity for more than 1 h per week had higher HRQOL measures in both physical and mental domains than those who were less physically active. Although many intervention trials have found a positive association between exercise and HRQOL, the available data from other intervention trials conducted among older adults is inconsistent. Additionally, information of the most effective mode of exercise that may influence HRQOL is lacking [32–35]. Self-efficacy is a possible psychological mediating factor and physical function is a possible physiological mediating factor. Previous research has shown that self-efficacy beliefs can be related to well-being following exercise interventions [36, 37] and that self-efficacy can explain adherence to exercise programs [38–41]. A central concept of the self-efficacy theory is so-called performance accomplishment, i.e. mastery experiences related to certain activities [42], and this points towards testing the mediating effect also of physical function.

The primary aim of this study is to explore the effects of a falls prevention programme, lasting 12 weeks, on HRQOL in older adults receiving home help services. Effects on the secondary outcomes, physical function and falls efficacy, will also be explored. A secondary aim of this study is to explore the mediating factors between falls prevention and HRQOL.

Methods

Study design

The study is a single-blinded, pragmatic RCT comparing one intervention group with a control group. The intervention group will receive an adapted version of the Otago Exercise Programme (OEP) over 12 weeks, while the control group receives usual care. Measurements are performed at baseline, at 3 months and at 6 months. The intervention and assessments will be conducted in the participants' homes. Assessors will be blinded to group participation.

Study setting and recruitment


Six municipalities in the Oslo region have agreed to take part in the research project. Participants are recruited through consultants in the municipalities coordinating and providing home help services. The researcher visits the municipalities on a regular basis to conduct the recruitment. Additionally, health workers in the municipalities are informed about the criteria to participate and will alert about eligible participants. Eligible participants will be contacted by the researcher by telephone and asked to consent to being sent information about the study. After a week, they will be contacted again to see if they consent orally to participate. Before baseline testing, the participants must provide a written informed consent. Figure 1 presents the planned flow of participants in the study.

Inclusion and exclusion criteria

Inclusion criteria are: Individuals who 1) are 67 or older, 2) receive home help services 3) have experienced at least one fall during the last 12 months, 4) are able to walk with or without a walking aid and 5) understand Norwegian. Exclusion criteria are: 1) medical contraindications to exercise, 2) life expectancy below 1 year, 3) a score below 23 on the Mini Mental State Examination (MMSE) and 4) currently participating in other falls prevention programmes or trials.

Randomisation

The participants are randomly assigned at a 1:1 ratio to the intervention group and the control group. A computer-generated, permuted block randomisation scheme is used to allocate the participants. Following randomisation, the participants receive information by



telephone on which group they are allocated to. See flow chart in Fig. 1.

In order to optimize the rigor of the RCT and to minimize bias, a number of methodological factors have been incorporated into the design of the study. The study participants will be randomly allocated to the groups via concealed allocation, as inadequately concealed allocation has been associated with bias in RCTs [43]. Due to the nature of the intervention, it is not possible to blind the participants or the treating therapists to the allocated groups. However, all assessors are blinded to the allocated groups. In further attempts to reduce bias, data will be analysed on an intention-to-treat basis. This preserves the randomisation process and imitate the real-life situation where the possibility exists that not all participants receive the prescribed treatment.

Study intervention

The intervention performed is based on the OEP, including home visits and motivational phone calls [44]. Balance exercises comprise tasks in standing, walking backwards, stair-walking and rising from a chair. Strengthening exercises uses ankle weight cuffs to strengthen hip extension and abduction, knee flexion and extension and ankle plantar and dorsiflexion. The programme also includes warm-up exercises as movement of neck and shoulders. The OEP has been described in more detail previously [44]. This programme has been shown to be effective in reducing number of falls and number of injuries resulting from falls in addition to improving strength and balance, and maintaining falls efficacy in home-dwelling older adults [45].

In previous studies the OEP has been performed over a period of 1 year [45]. A meta-analysis by Sherrington et al. [46], looking at the effect of falls prevention programmes, recommend a dose of at least 3 hours of exercise weekly for 6 months. This weekly dose is attempted, but the duration of 3 months is shorter than in previous studies. Nevertheless, as in the original OEP, the same number of home visits and telephone calls will be made, and the participants will be encouraged to do a sufficient amount of exercise between home visits. The rationale for the change in duration and frequency is both theoretical and pragmatic. Participants included for this study are frail older adults who have a fall history and who receive home help services. Previous research has shown that home help receivers fall more frequently and have a higher level of fear of falling [27, 47]. Poor health, fear of falling, depression and lack of strength are barriers for older adults in order to adhere to exercise programmes [48]. The participants in this study are thus likely to have a lower level of observance compared to more independent elderly and a duration of 1 year might be too long. Additionally, previous research has shown that falls prevention programmes which were considered too demanding by the participants even had a negative impact on QOL [49]. On the other hand, only receiving a few visits might not provide sufficient support which in turn could limit adherence. The pragmatic rationale relates to the organizational structure of physiotherapy services in the primary health care. For this group of older adults an intervention of 3 months is within the time frame of what the physiotherapists normally can provide. Finally, previous research has shown that also falls prevention programmes with a shorter duration than 6 months have had a positive effect on QOL [21].

The physiotherapists visit the participants at home five times during the intervention (week 1, 2, 4, 8 and 10) for instruction and for guiding the appropriate level and progression of each exercise. This includes one additional visit compared to the first 12 weeks of the original OEP intervention [44]. Each visit will take about 1 hour. The first visit may take longer as initial information is given and a relationship is established. At this visit advice related to safety when performing exercises is provided to the participants, both orally and written. In between supervised sessions, participants will be encouraged to continue exercising on their own three times weekly for 30 min. Equipment for exercising (ankle cuff weights of 1, 1.5, 2 and 2.5 kg) is provided for each participant. The weeks between home visits, the physiotherapists call the participants to motivate them to continue exercising and to answer possible questions. As a part of the programme, the participants are also encouraged to perform at least two or more weekly walks of ≥ 30 min. The participants are provided with a written exercise

booklet including illustrations. Following the intervention period, the participants to keep the exercise equipment and booklets, in order to continue exercising.

The participants in the control group will receive usual care from the primary health care service. Following re-assessment, the participants will have opportunity to participate in other falls prevention programmes, for example, already existing balance exercise group classes.

Education of intervention deliverers

Workshops and meetings will be held to inform the physiotherapists participating in the project. Before starting recruitment, one full day workshop on falls prevention and OEP is held for all therapists. Following start-up of recruitment and until the end of the project one workshop will be held approximately every fourth months. These last half a day and include one lecture on a topic concerning older people and time for discussion on the development of the project. Additional to the workshops, the researcher will have monthly meetings with the physiotherapists in the different municipalities. In order to make sure that the intervention is performed as intended, a fidelity checklist based on the OEP-manual has been developed. The physiotherapists use the checklist when conducting the home visits and phone calls.

Outcome measures

Following recruitment participants are assessed before they are randomised. Assessors are blinded to the participants' group assignment. The time window between baseline assessment and start of intervention is aimed to be within 2 weeks, and the same time window for assessments due at three and 6 months. Measurements and their order are selected to avoid physical and mental fatigue of the participants. Outcome measures that are employed have established reliability and validity, as recommended by the CONSORT group [50]. In addition to improving measurement quality and outcomes, it enables direct comparisons with other studies that investigate HRQOL and can possibly contribute to meta-analyses.

At baseline the Mini Mental State Examination (MMSE), a measurement of "Global cognitive function", is performed and is used as exclusion criteria. The maximum score is 30. A score below 23 indicates cognitive impairment and these participants are excluded [51]. Sociodemographic characteristics, like age and education, are also assessed at baseline. Primary and secondary outcome measures will be performed at baseline, at 3 months and at 6 months' follow-up.

Primary outcome variable

HRQOL is the primary outcome measured by the Short Form 36 Health Survey (SF-36) [52]. This is a generic

and validated questionnaire which, translated into Norwegian, is conducted as an interview [53]. The 36 items in SF-36 are grouped into eight health status scales: physical functioning, role limitations due to physical problems and due to emotional problems, bodily pain, general health perception, vitality, social functioning and mental health [52].

Secondary outcome variables

In addition to the SF36, the EQ-5D (1990 EuroQOL EQ-5D) is reported. The EQ-5D is a generic and validated questionnaire [54–57]. It describes five dimensions of HRQOL (mobility, self-care, usual activities, pain/discomfort, anxiety/depression), each of which can take one of five responses at five levels of severity (no problems/slight problems/moderate problems/severe problems/severe problems/extreme problems).

Physical function includes measures of balance, gait speed, muscle strength as well as activities of daily living. The Bergs Balance Scale is a 14-item scale, which is applied to assess static and dynamic balance in older adults [58]. Gait speed is assessed by measuring usual walking speed over four meters [59] and muscle strength is measured by the 30 s sit to stand test [60]. Instrumental ADL is recorded using the Norwegian Version of the Lawton IADL scale, which is a valid and reliable measure of a person's self-reported ability to perform complex activities of daily living [61].

Physical activity is measured using the "Walking habits questionnaire", a valid questionnaire for walking habits and physical activity for frailer older people [62]. This questionnaire assesses general behaviour of walking, regarding how often and for how long. The following questions are asked: "Do you take a daily walk?" (yes/no) or "If you do not take a daily walk how many times per week do you take a walk?" (never/almost never/1–2 days/3–4 days/ almost daily) and "How long does you walk generally last? (0–15 min/15–30 min/30–60 min/ 1 h–2 h/>2 h)". Walking time in minutes per week is calculated by taking the lowest level of days multiplied by lowest level of minutes for each response alternative [62].

Nutritional status is measured using the Mini Nutritional Assessment (MNA- elderly, Société des Produits Nestlé, S.A., Vevey, Switzerland) form. The first screening part of six questions is used which includes measurement of weight and height for calculation of BMI [63–65].

Falls efficacy is assessed using the Falls Efficacy Scale International (FES-I). This scale has shown good reliability and validity assessing concerns about falling in older adults, and is recommended for clinical trials and practise [66, 67]. It is a self-reported questionnaire, containing 16 items on different activities of daily living. Level

of concern is measured on a four-point scale ranging from 1, which is not at all concerned, to 4 which is very concerned [68].

Adherence to the programme is documented through an activity diary completed by the participants and a form checked by the physiotherapists during home visits and calls. Additionally, the participants have a falls calendar where they report adverse events. Adverse events are registered in the following four categories: falls, cardiovascular events, musculoskeletal injuries and health care utilization and will be documented as “due to the intervention” or “not due to the intervention” [69].

Sample size estimation

The sample size is estimated from the primary outcome, HRQOL (SF-36). A treatment difference of 10 points between the two groups in one of the domains in SF-36 is regarded to be of statistical and clinical significance. The associated standard deviation is assumed to be around 20 points. This implies a moderate effect size [70], which can be expected as previous OEP studies have shown substantial effects on physical outcomes [45]. Moreover, a similar Norwegian study, which included older adults performing exercise following discharge from hospital, estimated the required sample size identically [71]. Given a power of 80% and level $\alpha = 0.05$, we aim at including 150 participants, allowing for a 15–20% dropout, to detect a difference of 10 points between groups (see Fig. 1).

Statistical procedures

Statistical analysis is performed using SPSS or a similar statistical package. Descriptive data are reported for variables of interest. The data will be analysed following the intention to treat principle [72]. Prospective differences in primary and secondary outcomes and baseline characteristics between the intervention group and the control group will be assessed by t-tests for continuous and normal distributed variables and with non-parametric tests for categorical variables. Multiple linear regression modelling are used to control for confounding of between-group differences [73]. Hypotheses about mediating factors are tested through correlations, multiple regressions and bootstrapping methods exploring the correlations and explained variance of the chosen mediating factors and the changes in QOL [74]. Bootstrapping is a non-parametric method and is considered favourable with dichotomous variables (group 1 and 2) and small samples ($n < 250$) [74].

Discussion

The main purpose of the study is to evaluate the impact of OEP on HRQOL in older adults receiving home help services. We anticipate that the intervention described in this protocol will have a positive impact on the

HRQOL. The tailored intervention will have a potential to promote evidence-based decision-making and empower older people receiving home help services to remain to a greater extent in charge of their own lives. We rely on a systematic approach, which corresponds with the guidance on developing and evaluating RCTs [75]. Only a few studies have included HRQOL when measuring the effect of a falls prevention programme, and most of these studies include it as a secondary outcome [21, 76]. Outcomes examining HRQOL are selected based on literature identifying a standard set of measurements in falls prevention programmes [77, 78]. SF-36 is chosen due to its good validity, reliability and responsiveness when assessing older adults [79]. This outcome is detailed and broad, but it might be putting a burden on the participants due to its length and sensitive questions. Nevertheless, estimating HRQOL is important to determine whether the effect of a falls prevention programme is significant enough to achieve clinically relevant changes and thus to justify the implementation.

Several studies have looked at the effect of exercise on HRQOL, but to the authors' knowledge, none of them have specifically focused on older adults who receive home help services and who have a fall history. Studying the relationship between exercise and HRQOL is interesting due to the potential influence of exercise on both health and wellness through improvement of HRQOL [3, 80]. However, results from previous clinical exercise trials have reported mixed effects on HRQOL following exercise [32–35, 81]. Although many studies have found a positive association between exercise and HRQOL, available data from other trials is inconsistent and lacks information on the most effective mode of exercise that may influence HRQOL [32, 33, 71, 81]. This study can provide insight into the effect of falls preventative exercise and its applicability to home-dwelling older fallers with dependency of help from the primary health care service.

There are two ways an intervention mechanism can influence HRQOL, it can be a mediator or a moderator [82]. A mediating factor is defined as an intervening causal factor that may provide information concerning why the intervention increases HRQOL. Moderator mechanisms help us to understand for whom an intervention works [38] and can be classified as either characteristic of the person/group i.e. baseline characteristics or characteristics of the exercise protocol [38, 83]. Mediating mechanisms between HRQOL and falls-prevention programs may be both physiological, such as increased balance and strength, and psychological, such as self-efficacy [38]. A recent study provides evidence that fear is related to falls and concluded that falls self-efficacy plays a mediation role on the relationship between fear of falls and falls [84]. They recommend that any falls prevention should consider psychological covariates of falls,

especially subjects' self-efficacy to reduce falls, alongside other risk factors and covariates of falls. More theoretically driven research on these mechanisms behind treatment effects have been recommended [85–87].

It is widely accepted that falls and subsequent injuries are likely to result in a substantial reduction in quality of life for the persons affected as well a substantial economic burden to the healthcare system [88]. This provision of OEP in this setting could potentially be a beneficial and cost-effective intervention for this group of frail older adults, just as it is for community-dwelling older adults. Several studies have performed analysis on cost-effectiveness of exercise programmes which have shown that it can reduce healthcare costs [14, 89]. Due to its large sample size and theoretically based intervention the present study has the potential to generate new knowledge that may improve the design of future activity programmes for older fallers receiving home help services. Since both outcome measures as well as the intervention are carried out in a clinical setting, relevance and application of study findings to clinicians is enhanced. Results from this study will be primarily of interest to, and could be used by, health care managers and clinicians. Particularly, the results will be useful in decision making to set priorities relating to prevention measures in the community, to appropriately allocate resources and to assess costs and benefits of a falls prevention programme. Finally, the results can be useful for policy makers, in order to put preventative healthcare for this group of frail older adults on the agenda.

To conclude, older people receiving home help services represent a growing and diverse group as part of the population of community-dwelling older adults. The appropriate assessment of HRQOL, the mechanisms behind the relationship between fall prevention and HRQOL, the most effective mode of exercise, as well as the clinical relevance of the results, are challenging issues which are important to address.

Abbreviations

ADL: Activities of Daily Living; BMI: Body Mass Index; EQ-5D: European Quality of Life - 5 Dimensions; HRQOL: Health-related Quality of Life; iADL: Instrumental Activities of Daily Living; LEON: Lowest most efficient level of care; MMSE: Mini Mental State Examination; MNA: Mini Nutritional Assessment; OEP: Otago Exercise Programme; pADL: Personal Activities of Daily Living; QOL: Quality of Life; RCT: Randomised Controlled Trial; SF-36: Short Form 36 Health Survey; WHO: World Health Organization

Acknowledgements

Thanks to all physiotherapists and consultants working in the municipalities contributing with information on practical issues on recruitment and management of the intervention.

Funding

This study has received no external funding. It is internally funded by the Oslo and Akershus University College.

Availability of data and materials

The datasets generated and/or analysed during the proposed study are only available to the participating researchers due to data protection laws. Subsets or aggregation of these data will not include information that could compromise research participants' privacy. Consent can be made available from the corresponding author on reasonable request.

Authors' contributions

MB and AB were involved in choosing falls prevention programme as well as outcome measures. AB was responsible for the internal grant application for this trial. MB and AB contributed to the design of the study. MB administers the data collection and coordination of conducting the fall prevention programme. MB and AB wrote the first draft of the manuscript. TB and DAS critically revised and approved the final version of this manuscript.

Ethics approval and consent to participate

The project proposal has been approved by The Regional Committee for Medical Research Ethics in South Norway (Ref. 2014/2051). Informed consent is obtained from all participants included in the analyses, and the project is conducted according to the WMA Declaration of Helsinki.

Consent for publication

Not applicable

Competing interests

DAS is a Director of Later Life Training Ltd, a UK based non-profit organisation providing training to therapists in the effective delivery of the OEP to older adults.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Department of Physiotherapy, Faculty of Health Sciences, Oslo and Akershus University College, PO box 4 St. Olavs plass, Oslo 0130, Norway. ²Institute of Applied Health Research, School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, UK.

Received: 16 December 2016 Accepted: 7 August 2017

Published online: 14 August 2017

References

- Luthy C, Cedraschi C, Allaz A-F, Herrmann FR, Ludwig C. Health status and quality of life: results from a national survey in a community-dwelling sample of elderly people. *Qual Life Res.* 2015;24(7):1687–96.
- Salkeld G, Ameratunga SN, Cameron I, Cumming R, Easter S, Seymour J, Kurrle S, Quine S, Brown PM. Quality of life related to fear of falling and hip fracture in older women: a time trade off studyCommentary: older people's perspectives on life after hip fractures. *BMJ.* 2000;320(7231):341–6.
- Acree LS, Longfors J, Fjeldstad AS, Fjeldstad C, Schank B, Nickel KJ, Montgomery PS, Gardner AW. Physical activity is related to quality of life in older adults. *Health Qual Life Outcomes.* 2006;4(1):1.
- The WHOQOL Group. The World Health Organization quality of life assessment (WHOQOL): development and general psychometric properties. *Soc Sci Med.* 1998;46(12):1569–85.
- Helse- og omsorgsdepartementet. NOU 2011:11. Innovasjon i omsorg. Oslo: Departementenes servicesenter; 2011.
- Helse- og omsorgsdepartementet. St. Meld 47. Samhandlingsreformen. Oslo: Departementenes servicesenter; 2009.
- Christensen K, Doblhammer G, Rau R, Vaupel JW. Ageing populations: the challenges ahead. *Lancet.* 2009;374(9696):1196–208.
- Elkan R, Blair M, Robinson JJ. Evidence-based practice and health visiting: the need for theoretical underpinnings for evaluation. *J Adv Nurs.* 2000; 31(6):1316–23.
- Helse- og omsorgstjenesteloven. Lov om kommunale helse- og omsorgstjenester m.m. LOV-2011-06-24-30. 2011.
- Meinow B, Kåreholt I, Lagergren M. According to need? Predicting the amount of municipal home help allocated to elderly recipients in an urban area of Sweden. *Health Soc Care Community.* 2005;13(4):366–77.

11. Vikman I. Falls, perceived fall risk and activity curtailment among older people receiving home-help service. Doctoral thesis, Luleå: Luleå University of Technology; 2011.
12. Helsedirektoratet. Prioriteringer i helsesektoren. Verdigrunnlag, status og utfordringer. Oslo: Helsedirektoratet; 2012.
13. Vieira ER, Palmer RC, Chaves PH. Prevention of falls in older people living in the community. *BMJ*. 2016;353:i1419.
14. Gillespie LD, Robertson MC, Gillespie WJ, Sherrington C, Gates S, Clemson LM, Lamb SE. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev*. 2012;(9):Art. No.: CD007146. doi:10.1002/14651858.CD007146.pub3.
15. Bergland A. Falls suffered by the elderly living at home. Doctoral thesis, Oslo: University of Oslo; 2002.
16. Ambrose AF, Paul G, Hausdorff JM. Risk factors for falls among older adults: a review of the literature. *Maturitas*. 2013;75(1):51–61.
17. Vivanti AP, McDonald CK, Palmer MA, Sinnott M. Malnutrition associated with increased risk of frail mechanical falls among older people presenting to an emergency department. *Emerg Med Australas*. 2009; 21(5):386–94.
18. Saletti A, Johansson L, Yifter-Lindgren E, Wissing U, Österberg K, Cederholm T. Nutritional status and a 3-year follow-up in elderly receiving support at home. *Gerontology*. 2005;51(3):192–8.
19. Scheffer AC, Schuurmans MJ, Van Dijk N, Van Der Hooft T, De Rooij SE. Fear of falling: measurement strategy, prevalence, risk factors and consequences among older persons. *Age Ageing*. 2008;37(1):19–24.
20. Hawkins K, Musich S, Ozminkowski RJ, Bai M, Migliori RJ, Yeh CS. The burden of falling on the quality of life of adults with Medicare supplement insurance. *J Gerontol Nurs*. 2011;37(8):36–47.
21. Vaapio SS, Salminen MJ, Ojanlatva A, Kivelä S-L. Quality of life as an outcome of fall prevention interventions among the aged: a systematic review. *Eur J Public Health*. 2009;19(1):7–15.
22. Campbell AJ, Robertson MC. Rethinking individual and community fall prevention strategies: a meta-regression comparing single and multifactorial interventions. *Age Ageing*. 2007;36(6):656–62.
23. El-Khoury F, Cassou B, Charles MA, Dargent-Molina P. The effect of fall prevention exercise programmes on fall induced injuries in community dwelling older adults: systematic review and meta-analysis of randomised controlled trials. *BMJ*. 2013;347:f6234.
24. Petridou ET, Manti EG, Ntinapogias AG, Negri E, Szczerbńska K. What works better for community-dwelling older people at risk to fall? A meta-analysis of multifactorial versus physical exercise-alone interventions. *J Aging Health*. 2009; 21(5):713–29.
25. Sherrington C, Whitney JC, Lord SR, Herbert RD, Cumming RG, Close JC. Effective exercise for the prevention of falls: a systematic review and meta-analysis. *J Am Geriatr Soc*. 2008;56(12):2234–43.
26. Vikman I, Nordlund A, Näslund A, Nyberg L. Incidence and seasonality of falls amongst old people receiving home help services in a municipality in northern Sweden. *Int J Circumpolar Health*. 2011;70(2):195.
27. Meyer C, Dow B, Bilney BE, Moore KJ, Bingham AL, Hill KD. Falls in older people receiving in-home informal care across Victoria: influence on care recipients and caregivers. *Australas J Ageing*. 2012;31(1):6–12.
28. Fletcher PC, Hirdes JP. Restriction in activity associated with fear of falling among community-based seniors using home care services. *Age Ageing*. 2004;33(3):273–9.
29. Hellström Y, Persson G, Hallberg IR. Quality of life and symptoms among older people living at home. *J Adv Nurs*. 2004;48(6):584–93.
30. Markle-Reid M, Browne G, Gafni A, Roberts J, Weir R, Thabane L, Miles M, Vaitonis V, Hecimovich C, Baxter P. The effects and costs of a multifactorial and interdisciplinary team approach to falls prevention for older home care clients 'at risk' for falling: a randomized controlled trial. *Can J Aging*. 2010; 29(01):139–61.
31. Bonnefoy M, Boutitie F, Mercier C, Gueyffier F, Carre C, Guetemme G, Ravis B, Laville M, Cornu C. Efficacy of a home-based intervention programme on the physical activity level and functional ability of older people using domestic services: a randomised study. *J Nutr Health Aging*. 2012;16(4):370–7.
32. Chou C-H, Hwang C-L, Wu Y-T. Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: a meta-analysis. *Arch Phys Med Rehabil*. 2012;93(2):237–44.
33. Gillison FB, Skevington SM, Sato A, Standage M, Evangelidou S. The effects of exercise interventions on quality of life in clinical and healthy populations: a meta-analysis. *Soc Sci Med*. 2009;68(9):1700–10.
34. Kelley GA, Kelley KS, Hootman JM, Jones DL. Exercise and health-related quality of life in older community-dwelling adults: a meta-analysis of randomized controlled trials. *J Appl Gerontol*. 2009;28(3):369–94.
35. De Vries N, Van Ravensberg C, Hobbelen J, Rikkert MO, Staal J, Nijhuis-van der Sanden M. Effects of physical exercise therapy on mobility, physical functioning, physical activity and quality of life in community-dwelling older adults with impaired mobility, physical disability and/or multi-morbidity: a meta-analysis. *Ageing Res Rev*. 2012;11(1):136–49.
36. White SM, Wójcicki TR, McAuley E. Physical activity and quality of life in community dwelling older adults. *Health Qual Life Outcomes*. 2009;7(1):1.
37. McAuley E, Konopack JF, Morris KS, Motl RW, Hu L, Doerksen SE, Rosengren K. Physical activity and functional limitations in older women: influence of self-efficacy. *J Gerontol Ser B Psychol Sci Soc Sci*. 2006;61(5):P270–7.
38. Kraemer HC, Wilson GT, Fairburn CG, Agras WS. Mediators and moderators of treatment effects in randomized clinical trials. *Arch Gen Psychiatry*. 2002; 59(10):877–83.
39. McAuley E, Lox C, Duncan TE. Long-term maintenance of exercise, self-efficacy, and physiological change in older adults. *J Gerontol* 1993, 48(4): P218-P224.
40. McAuley E, Jerome GJ, Marquez DX, Elavsky S, Blissmer B. Exercise self-efficacy in older adults: social, affective, and behavioral influences. *Ann Behav Med*. 2003;25(1):1–7.
41. Landers DM, SM A. Physical activity and mental health. Hoboken, New Jersey: Wiley; 2007.
42. Bandura A. Self efficacy: the exercise of control. New York: W.H. Freeman; 1997.
43. Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias: dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA*. 1995;273(5):408–12.
44. Campbell AJ, Robertson MC. Otago Exercise programme to prevent falls in older adults. Wellington: ACC Thinksafe; 2003.
45. Robertson MC, Campbell AJ, Gardner MM, Devlin N. Preventing injuries in older people by preventing falls: a meta-analysis of individual-level data. *J Am Geriatr Soc*. 2002;50(5):905–11.
46. Sherrington C, Michaleff ZA, Fairhall N, Paul SS, Tiedemann A, Whitney J, Cumming RG, Herbert RD, Close JC, Lord SR. Exercise to prevent falls in older adults: an updated systematic review and meta-analysis. *Br J Sports Med*. Published Online First: October 4, 2016; doi:10.1136/bjsports-2016-096547.
47. Fletcher PC, Hirdes JP. Risk factors for falling among community-based seniors using home care services. *J Gerontol Ser A Biol Med Sci*. 2002;57(8): M504–10.
48. Forkan R, Pumper B, Smyth N, Wirkkala H, Ciol MA, Shumway-Cook A. Exercise adherence following physical therapy intervention in older adults with impaired balance. *Phys Ther*. 2006;86(3):401–10.
49. Vaapio S, Salminen M, Vahlberg T, Sjösten N, Isoaho R, Aarnio P, Kivelä S-L. Effects of risk-based multifactorial fall prevention on health-related quality of life among the community-dwelling aged: a randomized controlled trial. *Health Qual Life Outcomes*. 2007;5(1):1.
50. Schulz KF, Altman DG, Moher D. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMC Med*. 2010;8(1):1.
51. Tombaugh TN, McIntyre NJ. The mini-mental state examination: a comprehensive review. *J Am Geriatr Soc*. 1992;40(9):922–35.
52. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36): I. Conceptual framework and item selection. *Med Care*. 1992;473–83.
53. Loge JH, Kaasa S. Short form 36 (SF-36) health survey: normative data from the general Norwegian population. *Scand J Public Health*. 1998; 26(4):250–8.
54. The EuroQol Group. EuroQol—a new facility for the measurement of health-related quality of life. *Health policy*. 1990;16(3):199–208.
55. Brooks R, Group E. EuroQol: the current state of play. *Health Policy*. 1996;37(1):53–72.
56. Herdman M, Gudex C, Lloyd A, Janssen M, Kind P, Parkin D, Bonsel G, Badia X. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res*. 2011;20(10):1727–36.
57. Janssen M, Pickard AS, Golicki D, Gudex C, Niewada M, Scalone L, Swinburn P, Busschbach J. Measurement properties of the EQ-5D-5L compared to the EQ-5D-3L across eight patient groups: a multi-country study. *Qual Life Res*. 2013;22(7):1717–27.
58. Berg KO, Maki BE, Williams JI, Holliday PJ, Wood-Dauphinee SL. Clinical and laboratory measures of postural balance in an elderly population. *Arch Phys Med Rehabil*. 1992;73(11):1073–80.

59. Peters DM, Fritz SL, Krotish DE. Assessing the reliability and validity of a shorter walk test compared with the 10-meter walk test for measurements of gait speed in healthy, older adults. *J Geriatr Phys Ther.* 2013;36(1):24–30.
60. Jones CJ, Rikli RE, Beam WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exerc Sport.* 1999; 70(2):113–9.
61. Lawton M, Brody E. Assessment of older people: self-maintaining and instrumental activities of daily living. *Nurs Res.* 1970;19(3):278.
62. Frändin K, Grimby G, Mellström D, Svanborg A. Walking habits and health-related factors in a 70-year-old population. *Gerontology.* 1991;37(5):281–8.
63. Vellas B, Villars H, Abellan G, Soto M. Overview of the MNA® - its history and challenges. *J Nutr Health Aging.* 2006;10(6):456.
64. Rubenstein LZ, Harker JO, Salvà A, Guigoz Y, Vellas B. Screening for undernutrition in geriatric practice developing the short-form mini-nutritional assessment (MNA-SF). *J Gerontol Ser A Biol Med Sci.* 2001;56(6): M366–72.
65. Guigoz Y, Jensen G, Thomas D, Vellas B. The mini nutritional assessment (MNA®). Review of the literature - what does it tell us? *J Nutr Health Aging.* 2006;10(6):466.
66. Delbaere K, Close JC, Mikolaizak AS, Sachdev PS, Brodaty H, Lord SR. The falls efficacy scale international (FES-I). A comprehensive longitudinal validation study. *Age Ageing.* 2010;39(2):210–6.
67. Marques-Vieira CMA, Sousa LMM, Severino S, Sousa L, Caldeira S. Cross-cultural validation of the falls efficacy scale international in elderly: systematic literature review. *J Clinical Gerontol Geriatr.* 2016;7(3):72–6.
68. Yardley L, Beyer N, Hauer K, Kempen G, Piot-Ziegler C, Todd C. Development and initial validation of the falls efficacy scale-international (FES-I). *Age Ageing.* 2005;34(6):614–9.
69. Ory M, Resnick B, Jordan PJ, Coday M, Riebe D, Garber CE, Pruitt L, Bazzarre T. Screening, safety, and adverse events in physical activity interventions: collaborative experiences from the behavior change consortium. *Ann Behav Med.* 2005;29(2):20–8.
70. Cohen J. A power primer. *Psychol Bull.* 112(1):155–9.
71. Brovold T, Skelton DA, Bergland A. Older adults recently discharged from the hospital: effect of aerobic interval exercise on health-related quality of life, physical fitness, and physical activity. *J Am Geriatr Soc.* 2013;61(9):1580–5.
72. Hollis S, Campbell F. What is meant by intention to treat analysis? Survey of published randomised controlled trials. *BMJ.* 1999;319(7211):670–4.
73. Altman D. *Practical statistics for medical research.* London: Chapman and Hall; 1991.
74. Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods.* 2008;40(3):879–91.
75. Senn B, Kirsch M, Sanz C, Karlou C, Tulus K, De Leeuw J, Ringner A. Developing and evaluating complex interventions: the new Medical Research Council guidance. *Studies.* 2013;59:587–92.
76. McInnes E, Askie L. Evidence review on older people's views and experiences of falls prevention strategies. *Worldviews Evid Based Nurs.* 2004; 1(1):20–37.
77. Lamb SE, Jørstad-Stein EC, Hauer K, Becker C. Development of a common outcome data set for fall injury prevention trials: the prevention of falls network Europe consensus. *J Am Geriatr Soc.* 2005;53(9):1618–22.
78. Lamb SE, Becker C, Gillespie LD, Smith JL, Finnegan S, Potter R, Pfeiffer K. Reporting of complex interventions in clinical trials: development of a taxonomy to classify and describe fall-prevention interventions. *Trials.* 2011;12(1):1.
79. Haywood K, Garratt A, Fitzpatrick R. Quality of life in older people: a structured review of generic self-assessed health instruments. *Qual Life Res.* 2005;14(7):1651–68.
80. Vuillemin A, Boini S, Bertrais S, Tessier S, Oppert J-M, Hercberg S, Guillemin F, Briancçon S. Leisure time physical activity and health-related quality of life. *Prev Med.* 2005;41(2):562–9.
81. Heydarnejad S, Dehkordi AH. The effect of an exercise program on the health-quality of life in older adults. *Dan Med Bull.* 2010;57(4):113–7.
82. Michie S, Abraham C. Interventions to change health behaviours: evidence-based or evidence-inspired? *Psychol Health.* 2004;19(1):29–49.
83. Solberg PA. Exercise and well-being among older adults: A self-determination theory perspective. PhD thesis, Norwegian School of Sports Science. 2013.
84. Dadgari A, Hamid TA, Hakim MN, Mousavi SA, Dadvar L, Mohammadi M, Amerian N. The role of self-efficacy on fear of falls and fall among elderly community dwellers in Shahrud, Iran. *Nurs Pract Today.* 2016;2(3):112–20.
85. McAuley E, Blissmer B, Katula J, Duncan TE. Exercise environment, self-efficacy, and affective responses to acute exercise in older adults. *Psychol Health.* 2000;15(3):341–55.
86. Netz Y, Wu M-J, Becker BJ, Tenenbaum G. Physical activity and psychological well-being in advanced age: a meta-analysis of intervention studies. *Psychol Aging.* 2005;20(2):272.
87. Standage M, JL D. Motivational responses among older adults in sport and exercise settings. I Weiss MR (red) *Dev Sport Exerc Psychol lifespan Perspect.* 2004:357–81.
88. World Health Organization. WHO global report on falls prevention in older age. Geneva: World Health Organization; 2007.
89. Hektoen LF, Aas E, Lurås H. Cost-effectiveness in fall prevention for older women. *Scand J Public Health.* 2009;37(6):584–9.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

